





8–15 cm
3¼–6 in

ORANGE BIRCH BOLETE

Leccinum versipelle
F: Boletaceae

Found in Eurasia, this species has a yellow-orange fleshy cap. The stem has black, woolly flecks and the flesh stains lilac-black.



6–15 cm
2¼–6 in

BROWN BIRCH BOLETE

Leccinum scabrum
F: Boletaceae

One of many similar species, this fungus is from Eurasia and N. America. Its flesh may flush pink when cut, and the cap is sticky when wet.



1–2 cm
¾–¾ in

STALKED PUFFBALL-IN-ASPIC

Calostoma cinnabarinum
F: Calostomataceae

From N. America, this species' bright cinnabar-red ball on a stem emerges from a gelatinous layer.




5–100 cm
2–39 in

WET ROT

Coniophora puteana
F: Coniophoraceae

Found worldwide, this fungus forms a brown sheet of tissue, often warty or wrinkled, on wet timber. It causes serious damage to buildings.



5–9 cm
2–3½ in

BAROMETER EARTHSTAR

Astraeus hygrometricus
F: Diplocystidiaceae

Frequent in Eurasia and N. America, this species' star-like arms peel back to reveal the inner spore-filled ball, but close again in dry weather.



5–8 cm
2–3¼ in

CORNFLOWER BOLETE

Gyroporus cyanescens
F: Gyroporaceae

Growing on acidic soils in Eurasia and eastern N. America, this uncommon species is distinguished by its brittle, hollow stem.



1.5–5 cm
½–2 in

ROSY SPIKE

Gomphidius roseus
F: Gomphidiaceae

Growing in association with the bolete *Suillus bovinus* under pines in Eurasia, this fungus has a slimy, rose-pink cap with greyish gills.



4–8 cm
1½–3¼ in

COPPER SPIKE

Chroogomphus rutilus
F: Gomphidiaceae

Common under pines in Eurasia and western N. America, this species has a pointed, coppery brown cap.



6–15 cm
2¼–6 in

BROWN ROLLRIM

Paxillus involutus
F: Paxillaceae

Common in mixed woods in Eurasia and N. America, the downy, inrolled cap margin and soft yellow gills staining brown characterize this species.




4–10 cm
1½–4 in

COMMON EARTHBALL

Scloderma citrinum
F: Sclodermataceae

This potato-like fungus is common in damp woodland in Eurasia and N. America. Its thick skin has dark scales and a spore-filled black interior.



2–5 cm
¾–2 in

POTATO EARTHBALL

Scloderma bovista
F: Sclodermataceae

Common in woods in Eurasia and N. America, its smooth outer skin cracks into a fine mosaic. The inner purple-black spore-mass dries brownish.



2–8 cm
¾–3¼ in

FALSE CHANTERELLE

Hygrophoropsis aurantiaca
F: Hygrophoropsidaceae

Sometimes mistaken for a chanterelle, this species from Eurasia and N. America is distinguished by its crowded, multi-forked, soft gills.



5–10 cm
2–4 in

DYEBALL

Pisolithus arhizus
F: Sclodermataceae

This species is found worldwide on poor, sandy soils associated with pines. The egg-like spore-sacs in its interior are embedded in blackish jelly.



10–30 cm
4–12 in

VELVET ROLLRIM

Tapinella atrotomentosa
F: Tapinellaceae

Common on pine stumps in Eurasia and N. America, it has a cap with an inrolled margin; soft, thick gills; and a velvety stem.



4–10 cm
1½–4 in

WEEPING BOLETE

Suillus granulatus
F: Suillaceae

Common under pines in Eurasia and N. America, its ringless stem is covered with glandular dots.



5–10 cm
2–4 in

SLIPPERY JACK

Suillus luteus
F: Suillaceae

Found on pines in Eurasia and N. America, this fungus has a slimy cap and yellow pores. The large ring on its stem has a lavender underside.




3–7 cm
1¼–2¾ in

BOVINE BOLETE

Suillus bovinus
F: Suillaceae

Common under pines in Eurasia and N. America, this species has a sticky cap and angular, irregular pores.



5–10 cm
2–4 in

VELVET BOLETE

Suillus variegatus
F: Suillaceae

Frequent under pines in Eurasia, this species has a felty-scaly cap, with dark cinnamon-brown pores, and a ringless stem.



7–13 cm
2¾–5 in

LARCH BOLETE

Suillus grevillei
F: Suillaceae

Confined to larch woods in Eurasia and N. America, this species is yellow-orange to brick-red in colour. The stem has an ascending ring.