

CANTHARELLALES


The species of the Cantharellales order may look like agarics (members of the Agaricales order), but differ in several important respects. They may have fleshy fruitbodies with a cap and stem, but they lack true gills, having instead a smooth, wrinkled, or folded gill-like spore-producing surface on the underside. The spores are smooth and usually white to cream.



0.5–2 cm
3/16–3/4 in

HORN OF PLENTY
Craterellus cornucopioides
F: Cantharellaceae


A distinctive species found throughout Eurasia, this thin trumpet grows in troops in beech leaf litter. It produces a white spore deposit.



1–6 cm
3/8–2 1/4 in

TRUMPET CHANTERELLE
Craterellus tubaeformis
F: Cantharellaceae

Often found in large troops in mixed woods in Eurasia and N. America, this fungus has a variable colour and shallow, blunt veins rather than gills.



irregular, often lobed cap

5–15 cm
2–6 in

WOOD HEDGEHOG
Hydnum repandum
F: Hydnaceae

Occurring in Eurasia and N. America, this dusky orange species is irregular in shape, with tiny spines underneath the cap.



cap often depressed at centre


2–12 cm
3/4–4 3/4 in

CHANTERELLE
Cantharellus cibarius
F: Cantharellaceae

Found in Eurasia and N. America, this species has blunt-edged gills with numerous cross-veins and a smell of apricots.

inrolled margin

tapered stem



3–8 cm
1 1/4–3 1/4 in

CRESTED CORAL
Clavulina coralloides
F: Clavulinaceae

Very common in woods in Eurasia and N. America, this fungus forms a coral-like mass of white branches, each dividing into fine points.

GEASTRALES


The Geastrales, or earthstars, share the common character of a thick outer layer – the peridium – which splits apart and peels back to form star-like arms. These reveal a central spore-sac, like a puffball, from which the dark brown, warted spores exit via a pore at the apex. Earthstars are found in leaf litter and bare, sandy soils – for example, in sand dunes.



3–6 cm
1 1/4–2 1/4 in

SESSILE EARTHSTAR
Geastrum fimbriatum
F: Geastraceae

Found in Eurasia and N. America, this fungus's globe-shaped, pale brown fruitbody splits open with 5–9 arms. Its greyish spore-sac has a fringed opening.



5–8 cm
2–3 1/4 in

ARCHED EARTHSTAR
Geastrum fornicatum
F: Geastraceae

This earthstar's arms stay attached to a disc of tissue on the soil, and its stalked spore-sac has a prominent opening. It occurs in Eurasia and N. America.

STRIATE EARTHSTAR

Geastrum striatum
F: Geastraceae

Found in Eurasia, this is one of the smaller earthstars. Its pale grey spore-sac is stalked and has a sharply pointed, striated opening.

3–6.5 cm
1 1/4–2 1/2 in



collar below spore-sac



4–12 cm
1 1/2–4 3/4 in

COLLARED EARTHSTAR
Geastrum triplex
F: Geastraceae

Growing in Eurasia and N. America, this is the most common earthstar. Its rays crack to leave a cup-like collar around the spore-sac.



7–15 cm
2 3/4–6 in

PEPPER POT
Myriostoma coliforme
F: Geastraceae

Found on dry, sandy soils in Eurasia and N. America, this rare species has a distinctive, large spore-sac with several pore openings.