

GOMPHALES

Although some species were included with the chanterelles in Cantharellales, DNA analysis of the Gomphales fungi suggests that they are more closely related to the stinkhorn fungi in the Phallales. They often form large fruitbodies, varying in shape from simple clubs (in genus *Clavariadelphus*) to trumpet- or chanterelle-like structures with a complex spore-producing surface (in genus *Gomphus*).



GIANT CLUB

Clavariadelphus pistillaris
F: Clavariadelphaceae

A rare species of Eurasia and N. America, this fungus forms a large, swollen club, with a smooth to slightly wrinkled surface, which bruises purple-brown.



SCALY VASE CHANTERELLE

Gomphus floccosus
F: Gomphaceae

Common in N. America, this fungus resembles a fleshy, trumpet-shaped vase with scales at the top and wrinkled "gills" on its undersurface.


stains green with age



GREENING CORAL

Phaeoclavulina abietina
F: Gomphaceae

Found in conifer woods in Eurasia and N. America, this yellow-olive fungus has densely packed branches that bruise green.



UPRIGHT CORAL


Ramaria stricta
F: Gomphaceae

A fairly common species in Eurasia and N. America, it is always attached to decaying wood or woodchip mulch. The branches are pale brown, bruising reddish.

ROSSO CORAL

Ramaria botrytis
F: Gomphaceae

Found in beech woods in Eurasia and N. America, this uncommon coral fungus has pinkish white branches with deep red tips.



GLOEOPHYLLALES

This is an order of wood-decay fungi that is characterized by the ability to produce a brown rot of wood. The order Gloeophyllales has a single family, the Gloeophyllaceae, which includes the genus *Gloeophyllum*. Some well-known bracket fungi on conifer timber are members of this genus.




ANISE MAZEGILL

Gloeophyllum odoratum
F: Gloeophyllaceae

This fungus, which grows on decayed conifer wood, is found in Eurasia and N. America. It has irregular brackets with yellow pores and an aniseed odour.

DACRYMYCETALES

This order forms very simple, rounded or branched gelatinous fruitbodies, usually with bright orange coloration. Smooth or wrinkled, the Dacrymycetales have unusual spore-producing cells (basidia), which typically have two stout stalks (sterigmata), each bearing a spore. They mainly feed on dead wood.



YELLOW STAGSHORN

Calocera viscosa
F: Dacrymycetaceae

Attached to conifer wood, this fungus is found in Eurasia and N. America. Its clubs usually divide into branches with a gelatinous, rubbery texture.

HYMENOGASTRALES

This group contains a number of diverse types of fungi including some crust fungi, polypores, such as *Mensularia* and *Phellinus*, as well as several agaric species, such as *Rickenella*. The Hymenochaetales are defined through molecular studies and have few uniting physical characteristics. Many feed on wood and may cause a white rot of timber.



thick, pale margin

ALDER BRACKET


Mensularia radiata
F: Hymenochaetaceae

This deep reddish brown fungus with a paler margin often forms vertical chains of brackets on alder and other trees in Eurasia and N. America.

TIGER'S EYE

Coltricia perennis
F: Hymenochaetaceae

Frequently found in acidic heathlands of Eurasia and N. America, this fungus's goblet-shaped, thin fruitbodies are concentrically zoned.



OAK CURTAIN CRUST

Hymenochaete rubiginosa
F: Hymenochaetaceae

Found across Eurasia, this fungus forms overlapping brackets and crusts mainly on fallen oak timber. Its tough fruitbodies have concentric markings.

WILLOW BRACKET

Phellinus igniarius
F: Hymenochaetaceae

This grey to almost black perennial bracket from Eurasia and N. America grows over many years. It is hoof-shaped and extremely woody.



ORANGE MOSSCAP

Rickenella fibula
F: Incertae sedis in Rickenella clade

This tiny species is common in mossy grasslands of Eurasia and N. America. Its bright orange cap is radially marked with a darker centre.

