



5–30 cm
2–12 in

TINDER BRACKET
Fomes fomentarius
F: Polyporaceae

This grey-brown, hoof-shaped, perennial bracket grows on birches and other deciduous trees. It is found in Eurasia and N. America.

10–60 cm
4–23½ in

DRYAD'S SADDLE
Cerioporus squamosus
F: Polyporaceae

Found in early summer in Eurasia and N. America, these circular or fan-shaped brackets have concentric scales and pores on the underside.



8–15 cm
3¼–6 in

BLUSHING BRACKET
Daedaleopsis confragosa
F: Polyporaceae

One of the commonest brackets, especially on willows, in Eurasia and N. America, this semicircular species has cream pores that bruise pinkish red.



UPPER SURFACE

3–10 cm
1¼–4 in

BIRCH MAZEGILL
Lenzites betulina
F: Polyporaceae

Usually found on birches in Eurasia and N. America, this tough, leathery species has pores that can be extremely elongated in gill-like ridges.



UNDERSIDE



5–20 cm
2–8 in

velvety
stem base

BLACK FOOTED POLYPORE
Polyporus durus
F: Polyporaceae

This fungus has a funnel-shaped, leathery cap and its stem is black at the base. It grows on fallen beech logs in Eurasia and N. America.



5–20 cm
2–8 in

TUBEROUS POLYPORE
Polyporus tuberaster
F: Polyporaceae

Growing on fallen branches in Eurasia and N. America, this species may root into the ground and form a large, tuberous mass.



2–4 cm
¾–1½ in



3–8 cm
1¼–3¼ in

WINTER POLYPORE
Lentinus brumalis
F: Polyporaceae

This small species grows on fallen branches in Eurasia and N. America. It has rather large, decurrent (running down the stem) pores. The stem is central or off-centre.



3–10 cm
1¼–4 in

CINNABAR BRACKET
Pycnoporus cinnabarinus
F: Polyporaceae

Growing on dead deciduous timber in Eurasia and N. America, this rare species has a bright reddish orange, leathery, annual bracket.

PURPLEPORE BRACKET
Trichaptum abietinum
F: Polyporaceae

Found on fallen conifers in Eurasia and N. America, this fan-shaped bracket has concentric zones of pale grey, often tinged green by algae, with a purple margin.



5–12 cm
2–4¾ in

HAIRY BRACKET
Trametes hirsuta
F: Polyporaceae

This semicircular bracket is covered in minute hairs. It grows on dead deciduous wood or sometimes on old gorse stems in Eurasia and N. America.



TURKEYTAIL
Trametes versicolor
F: Polyporaceae


Found in Eurasia and N. America in a variety of colours, this species has brackets with concentric zones of different colours and a white pore surface.

2–7 cm
¾–2¾ in

LUMPY BRACKET
Trametes gibbosa
F: Polyporaceae

This cream-coloured bracket is often stained green from algae. Its pores may be long and maze-like. It grows on fallen deciduous logs in Eurasia and N. America.

10–30 cm
4–12 in




small-lobed
bracket

2–6 cm
¾–2¼ in

HEN OF THE WOODS
Grifola frondosa
F: Sparassidaceae

Found in Eurasia and N. America, this species grows at the base of oak, often where lightning has struck. It has a dense cluster of small brackets.



basal stem



fleshy, bright
yellow-orange
bracket

10–40 cm
4–16 in

WOOD CAULIFLOWER
Sparassis crispa
F: Sparassidaceae

Growing at the base of conifers in Eurasia and N. America, this species has cream lobes that are flattened and fleshy like a cauliflower.

