




dry, smooth
cap surface

5–15 cm
2–6 in

CHARCOAL BURNER
Russula cyanoxantha
F: Russulaceae

Growing in mixed woods in Eurasia and N. America, this mushroom's cap varies from purple-lilac to uniformly green. The gills are forked, flexible, and greasy.



5–10 cm
2–4 in

BLOODY BRITTLEGILL
Russula sanguinaria
F: Russulaceae


Associated with pines in Eurasia and N. America, this species has a scarlet cap and red-streaked stem. Its spores are pale ochre.



3–7 cm
1¼–2¾ in

BEECHWOOD SICKENER
Russula mairei
F: Russulaceae


Exclusively found with beech in Eurasia, this species has a scarlet cap and bluish white gills.



4–9 cm
1½–3½ in

GREEN BRITTLEGILL
Russula aeruginea
F: Russulaceae

Common under birches in Eurasia and N. America, this mushroom's pale olive to grass-green cap has tiny rust-coloured spots. Its spores are pale cream.



5–10 cm
2–4 in

YELLOW SWAMP BRITTLEGILL
Russula claroflava
F: Russulaceae

A common species on moss in boggy birch woods in Eurasia and N. America, its cap, yellowish cream gills, and white stem all bruise grey-black.



5–12 cm
2–4¾ in

OGHRE BRITTLEGILL
Russula ochroleuca
F: Russulaceae

One of the commonest species found in Eurasia, it is identifiable by its matt yellow-ochre or greenish yellow cap and white gills.



4–10 cm
1½–4 in

PRIMROSE BRITTLEGILL
Russula sardonia
F: Russulaceae

Found in Eurasia, this species grows in association with pine. Variable in colour, from purple to green or yellow, it has a fruity odour.




3–8 cm
1¼–3¼ in

SICKENER
Russula emetica
F: Russulaceae

Growing in wet pine woods in Eurasia and N. America, the sickener has a bright scarlet cap that contrasts with its pure white gills and stem.

dry and matt
cap surface

gills may have
red edges



4–12 cm
1½–4¾ in

ROSY BRITTLEGILL
Russula lepida
F: Russulaceae

Found in Eurasia, this fungus has a carmine-red, dry, hard cap that fades rapidly. Its stem may also be red and the flesh has a smell similar to cedar wood.

stem often
flushes red



6–15 cm
2¼–6 in

CRAB BRITTLEGILL
Russula xerampelina
F: Russulaceae

Found in Eurasia and N. America, this fungus is one of many related species that can only be identified under a microscope or by habitat.



8–15 cm
3¼–6 in

STINKING BRITTLEGILL
Russula foetens
F: Russulaceae

This large, orange-brown fungus with a grooved, lumpy cap margin grows in Eurasia and N. America. It has a sour, rancid odour.



2–6 cm
¾–2¼ in

HAIRY CURTAIN CRUST
Stereum hirsutum
F: Stereaceae

Found in Eurasia and N. America, this mushroom varies in shape from fully crust-like to small overlapping brackets. It is hairy above and smooth below.



10–50 cm
4–20 in

BLEEDING BROADLEAF CRUST
Stereum rugosum
F: Stereaceae

Growing in Eurasia and N. America, this species forms small crusts, and occasionally small brackets on wood. Its upper surface bleeds red where cut.



0.5–2 cm
¾–¾ in

EARPICK FUNGUS
Auriscalpium vulgare
F: Auriscalpiaceae

Growing on pine cones in Eurasia and N. America, this unique mushroom looks like a bent-over spoon with minute spines hanging down from the small, furry cap.



5–25 cm
2–10 in

ROOT ROT
Heterobasidion annosum
F: Bondarzewiaceae

Usually parasitic on conifers in Eurasia and N. America, this species has a pale brown crust that darkens with age.



10–40 cm
4–16 in

CORAL TOOTH
Hericium coralloides
F: Hericiaceae

Usually found on beech trees in Eurasia and N. America, this endangered species has white fruitbody branches with pendent spines on the lower surfaces.