


HELOTIALES

The fungi of this order are distinguished by their disc- or cup-shaped fruitbodies, unlike some similar cup fungi. Their sac-like spore-producing cells, or asci, do not have an apical lid (a flap through which they open). Most members live on humus-rich soil, dead logs, and other organic matter. This order also includes some of the most damaging plant parasites.



TROCHILA ILICINA
F: Dermateaceae

Abundant on fallen holly leaves, this fungus is found in Eurasia and N. America. It forms spore-producing spots on the leaf surface.



ROSE BLACK SPOT
Diplocarpon rosae
F: Dermateaceae

Commonly found on rose leaves in Eurasia and N. America, this fungus causes black spots, which merge together to cause large black patches.



merged black spots

spore-producing inner surface

smooth surface of cup

ANEMONE CUP
Dumontinia tuberosa
F: Sclerotiniaceae

Parasitic on the tubers of anemones, this common Eurasian species has a long, black stem and a simple brown cup.

SCALY EARTHTONGUE
Geoglossum fallax
F: Geoglossaceae


Growing in meadows in Eurasia and N. America, this uncommon fungus is one of several species with blackish, flattened clubs, which can only be separated using a microscope.



0.5–4 cm
3/16–1 1/2 in

BLACK BULGAR
Bulgaria inquinans
F: Bulgariaceae

Found in Eurasia and N. America, this species has a brown outer surface. Its spore-producing inner surface is smooth, black, and rubbery.



APPLE BROWN ROT
Monilia fructigena
F: Sclerotiniaceae

Very common in Eurasia, this fungus occurs primarily on apples and pears, although it is also found on *Prunus* sp., causing brown fruit rot.

0.3–1 cm
1/8–3/8 in

JELLY BABIES

Leotia lubrica
F: Leotiaceae

Frequently found in mixed woodland in Eurasia and N. America, this species has a lobed head with a margin that rolls back on itself.



GREEN ELF CUP
Chlorociboria aeruginascens
F: Helotiaceae

Found in Eurasia and N. America, the blue-green mature cups are rather uncommon, but the green stains they cause on fallen oak logs are easily visible.



0.5–1.5 cm
3/16–1/2 in

BROWN CUP
Rutstroemia firma
F: Rutstroemiaceae

Consisting of a light brown cup and a narrow stem, this fungus grows on fallen branches, especially of oaks, in Europe. It turns the host wood black.



BEECH JELLYDISC
Neobulgaria pura
F: Helotiaceae

Frequently found on fallen beech logs in Eurasia, this translucent jellydisc varies from pale pink to pale lilac. The discs are often distorted from crowding together.

0.5–3 cm
3/16–1 1/4 in



1–3 mm
1/32–1/8 in

LEMON DISCO
Bispora citrina
F: Helotiaceae

Very common on dead hardwood in Eurasia, this species occurs in clusters. Its golden-yellow discs sometimes cover entire branches.



0.5–2 cm
3/16–3/4 in

LARGE PURPLE DROP
Ascocoryne cylichnium
F: Helotiaceae

Quite common on fallen beech logs in Eurasia, this species attaches itself centrally to the wood and forms gelatinous, irregular discs when sexually mature.



0.2–1 cm
1/16–3/8 in

BOG BEACON
Mitrula paludosa
F: Helotiaceae

This species, found in Eurasia and N. America, grows on plant remains in shallow water in spring and early summer. It has a rounded to tongue-shaped head.