

PEZIZALES

Members of this order produce spores inside sac-like structures, or asci, which typically open by rupturing to form an operculum (terminal lid) and eject the spores. The order includes a number of species of economic importance, such as morels, truffles, and desert truffles.




COMMON EARTHCUP
Geopora arenicola
F: Pyronemataceae

Common in Eurasia, this fungus is difficult to spot as it lies buried in sandy soils. It has a smooth, spore-producing inner surface.

HARE'S EAR
Otidea onotica
F: Pyronemataceae

This common species is often found in clusters in broadleaf forests in Eurasia and N. America. Its tall cups are split down on one side.



COMMON EYELASH
Scutellinia scutellata
F: Pyronemataceae

This is one of many similar species; its fruitbody is a scarlet cup with a fringe of black hairs. This fungus is common on wet, rotten wood in Eurasia and N. America.



TOOTHED CUP
Tarzetta cupularis
F: Pyronemataceae

A common species in alkaline soils in woods of Eurasia and N. America, its goblet-like cup has a short stem.




FALSE MOREL
Gyromitra esculenta
F: Discinaceae

This poisonous species is found throughout N. America and Eurasia, and usually grows under conifers in spring. The shiny brown cap looks like a wrinkled brain.

ORANGE PEEL FUNGUS

Aleuria aurantia
F: Pyronemataceae

Often found along gravelly dirt tracks in Eurasia and N. America, this is an unmistakable species, with thin, orange caps.



BLEACH CUP
Disciotis venosa
F: Morchellaceae

Growing in spring in damp woodland in Eurasia and N. America, this short-stemmed species has a chlorine-like smell. Its inner surface is brown and wrinkled and the outer surface is pale.

HALF-FREE MOREL
Morchella semilibera
F: Morchellaceae

This hollow morel looks like a dark, ridged thimble on a scurfy, pale stem. It is common in spring in mixed woods in Eurasia and N. America.

BLACK MOREL
Morchella elata
F: Morchellaceae

Common in woods during spring in Eurasia and N. America, this species has a pinkish buff to black cap, with cross-connected black ridges, and a hollow stem.




MOREL
Morchella esculenta
F: Morchellaceae

This prized species grows in spring in calcareous woodland in N. America and Eurasia. It has a sponge-like, hollow cap and a hollow stem.

THIMBLE MOREL
Verpa conica
F: Morchellaceae

An uncommon species, this fungus grows in woods and hedges on chalky soils in Eurasia and N. America. Its smooth, thimble-like cap sits on a hollow stem.



BLISTERED CUP
Peziza vesiculosa
F: Pezizaceae

Common in Eurasia and N. America, this fungus typically occurs in clusters on compost, straw, or manure. Its brittle cups can have ragged edges.



CELLAR CUP
Peziza cerea
F: Pezizaceae

Found across Eurasia and N. America, this fungus often occurs on damp brickwork. It is dark ochre on the inside and pale outside.



BAY CUP
Peziza badia
F: Pezizaceae

One of many similar species, this common fungus of Eurasian and N. American woods has a spore-producing inner surface. Its brown cup develops an olive tint with age.

