

» LICHENS



ORANGE LICHEN
Caloplaca verruculifera
F: Teloschistaceae

This lichen has radiating lobes with apothecia (spore-producing discs) at the centre. It is found on rocks in coastal areas in Eurasia and N. America, often near bird perches.

5–10 cm
2–4 in



GOLDEN-EYE LICHEN
Teloschistes chrysophthalmus
F: Teloschistaceae

A critically endangered species in Eurasia, America, and the tropics, this lichen grows on shrubs and small trees in old orchards and hedgerows. Its branched lobes produce large, orange discs.


2.5–7.5 cm
1–3 in

2.5–7.5 cm
1–3 in

COMMON WALL LICHEN
Xanthoria parietina
F: Teloschistaceae

Growing on trees, walls, and roofs in N. America, Eurasia, Africa, and Australia, this lichen has rounded, yellow-orange lobes.

rounded lobes
that lift at edges



5–15 cm/2–6 in



BEARD LICHEN
Usnea filipendula
F: Parmeliaceae

Found mainly in northern regions, this species forms green-grey, pendent clumps on trees. Spiny apothecia form at the tips.



2.5–7.5 cm
1–3 in

REINDEER MOSS
Flavocetraria nivalis
F: Parmeliaceae

Found on mountain heaths and upland moors in N. America and Eurasia, this species has brownish, flattened leafy fronds with spiny margins.



2.5–7.5 cm
1–3 in

HAMMERED SHIELD LICHEN
Parmelia sulcata
F: Parmeliaceae


This lichen's flattened lobes are grey-green, with rounded tips and powdery, reproductive structures on the surface. It is commonly found on trees in N. America and Eurasia.



2.5–7.5 cm
1–3 in

HOARY ROSETTE LICHEN
Physcia aipolia
F: Physciaceae

Growing on tree bark in Eurasia and the Americas, this grey to brownish grey species forms rough patches with a lobed margin. It has black apothecia.



2.5–7.5 cm
1–3 in

2.5–7.5 cm
1–3 in


POWDER-HEADED TUBE-LICHEN
Hypogymnia tubulosa
F: Parmeliaceae

Common on twigs and tree trunks, this lichen has lobes that are grey-green above and dark underneath. It is found in Eurasia and N. America.



HOODED TUBE-LICHEN
Hypogymnia physodes
F: Parmeliaceae

Found worldwide on trees, rocks, and walls, this lichen has pale grey-green lobes with wavy margins. The rare apothecia are red-brown with grey margins.



1–5 cm
½–2 in

DEVIL'S MATCHSTICK
Cladonia floerkeana
F: Cladoniaceae


This species is common on peaty soil in Eurasia and N. America. It forms a crust of greenish grey scales from which stalks emerge, tipped with scarlet apothecia.



2.5–12.5 cm
1–5 in

CARTILAGE LICHEN
Ramalina fraxinea
F: Ramalinaceae


Growing on trees in Eurasia and N. America, this species forms flattened, grey-green branches dotted with apothecia.



2.5–10 cm
1–4 in

CORAL LICHEN
Sphaerophorus globosus
F: Sphaerophoraceae

Found on rocks in mountainous areas of N. Eurasia and N. America, this fungus forms dense cushions of pinkish brown branches with globular apothecia.



2.5–10 cm
1–4 in

BLACK SHIELDS
Tephromela atra
F: Mycoblastaceae


This lichen has pale grey crustose lobes that look like dried porridge. Its apothecia are black. The lichen is found on exposed rocks in N. America and Eurasia.



2.5–10 cm/1–4 in

STONEWALL RIM-LICHEN
Lecanora muralis
F: Lecanoraceae

This lichen often grows on concrete and on rocks. Its grey-green lobes radiate outwards. It is found in Eurasia and N. America.



2.5–10 cm
1–4 in

REINDEER LICHEN
Cladonia portentosa
F: Cladoniaceae

One of several reindeer lichens, this species is common on heath and moorland in N. America and Eurasia. Its thin, hollow branches repeatedly divide.