



10–20 cm
4–8 in

FLOWERPOT CORAL
Goniopora columna
F: Poritidae

This species has daisy-like polyps that are greatly lengthened when fully extended. It is an Indo-Pacific relative of the lobe coral.

4–5 m
13–16 ft



LOBE CORAL
Porites lobata
F: Poritidae

One of the commonest reef-builders of the Indo-Pacific, this coral forms large encrusting colonies in places buffeted by strong wave action.


1–2 m
3¼–6½ ft



LOBED BRAIN CORAL
Lobophyllia sp.
F: Lobophylliidae

Massive colonies of this brain coral – either flattened or dome-shaped – occur on reefs in tropical waters of the Indo-Pacific.

10–12 cm
4–4¾ in



DAHLIA ANEMONE
Urticina felina
F: Actiniidae

Sticky swellings on this anemone may carry so much debris that, with tentacles retracted, it resembles a small pile of gravel. It occurs around the North Pole.

1–3 m
3¼–9¾ ft



STAGHORN CORAL
Acropora sp.
F: Acroporidae

The branching stag horn corals are among the largest tropical reef-builders. Nourished by photosynthetic algae, these species are fast growing.



1 m
3¼ ft

DAISY CORAL
Goniopora sp.
F: Poritidae

Long-reaching polyps of *Goniopora* stony corals on tropical reefs typically have 24 tentacles. They are among the most flower-like of corals.


1–3 m
3¼–10 ft



LARGE-GROOVED BRAIN CORAL
Colpophyllia sp.
F: Faviidae


The hemispherical brain-like structure of this coral is typical of its family. It is a tropical reef-builder that contains photosynthetic algae.

10–20 cm
4–8 in



MUSHROOM CORAL
Fungia fungites
F: Fungiidae

This is one of many tropical mushroom corals that are not reef-building, but live among other species as solitary polyps crawling over the ocean floor.



50–100 cm
20–40 in

MAGNIFICENT SEA ANEMONE
Heteractis magnifica
F: Stichodactylida

This giant anemone is from Indo-Pacific reefs, where it lives in close partnership with a variety of fish species, including clown fish.

2.5–15 cm
1–6 in



PLUMOSE ANEMONE
Metridium senile
F: Metridiidae

This is a globally widespread member of a family of anemones that is characterized by a fuzzy mass of tentacles. It can split to form genetically identical populations.

5–7 cm
2–2¾ in



SNAKELOCKS ANEMONE
Anemonia viridis
F: Actiniidae

The conspicuously long tentacles of this common European intertidal anemone rarely retract, even when exposed at low tide.


100–200 m/300–600 ft



ATLANTIC COLD WATER CORAL
Desmophyllum pertusum
F: Caryophylliidae

Unlike many deep sea corals lacking nutrient-supplying algae, this North Atlantic species forms extensive reefs, although only very slowly.


10–15 mm
¾–½ in



DEVONSHIRE CUP CORAL
Caryophyllia (Caryophyllia) smithii
F: Caryophylliidae

This N. E. Atlantic species belongs to a family containing cold water corals, some with large, anemone-like polyps. Barnacles frequently attach to it.


10–15 cm
4–6 in



COMMON TUBE ANEMONE
Cerianthus membranaceus
F: Cerianthidae

Tube anemones burrow through sediment, using unique non-stinging cnidocytes to build felt-like tubes from mucus. This species occurs on European offshore mud.

50–100 cm
20–40 in



ATLANTIC BLACK CORAL
Antipathes sp.
F: Antipathidae

Mostly found in deep waters, black corals consist of spiny polyp colonies encased in slender, horny exoskeletons.