


**AMERICAN SUN SPIDER***Eremobates* sp.


F: Eremobatidae

Eremobates are sun spiders with particularly large jaws. This nocturnal species is found in warm parts of N. and C. America.

**PAINTED SUN SPIDER***Metasolpuga picta*

F: Solpugidae

Sun spiders, also called camel spiders, are fast-running, desert-living relatives of true spiders. This is a diurnal species from Namibia.

**WIND SPIDER***Galeodes arabs*

F: Galeodidae

A common species in the largest genus of sun spiders, this Middle Eastern species is named for its ability to tolerate sand storms.

**FLOUR MITE***Acarus siro*

F: Acaridae

This is an important pest that feeds on stored cereal products. Like many other species, its presence can cause allergic reactions in humans.

**AUTUMN CHIGGER***Neotrombicula autumnalis*


F: Trombiculidae

Adults of chigger mites are vegetarian, but their larvae feed on the skin of other animals, including humans. Their chewing causes intense irritation.

**COMMON VELVET MITE***Trombidium holosericeum*


F: Trombiculidae

This is a widespread Eurasian species of velvet mite. When young, it lives parasitically off other arthropods, but becomes predatory when mature.

**RED SPIDER MITE***Tetranychus urticae*

F: Tetranychidae

Spider mites are a family of mites that suck the sap of plants. They weaken the plant and can transmit viral diseases.

**LONE STAR TICK***Amblyomma americanum*

F: Ixodidae

Like other blood-sucking ticks, this species – common in woodlands in the USA – can transmit a number of disease-causing microbes.

**VARROA MITE***Varroa* sp.


F: Varroidae

Parasitic on honeybees, young of this mite feed on bee larvae. When mature, they attach to and feed on adult bees and spread to other hives.

**MANGE MITE***Sarcoptes scabiei*

F: Sarcoptidae

This tiny mite burrows into the skin of various mammal species, completing its life cycle there. It causes scabies in humans and mange in carnivores.

**CHICKEN MITE***Dermanyssus gallinae*


F: Dermanyssidae

A blood-sucking parasite of poultry, this mite lives – and completes its life cycle – in crevices away from its host, but emerges at night to feed.

**PERSIAN FOWL TICK***Argas persicus*

F: Argasidae

A blood-sucking parasite of fowl, including domestic chickens, this oval-shaped, soft-bodied tick can spread disease between birds and can cause paralysis.

**SPINY HARVESTMAN***Discocyrtus* sp.

F: Gonyleptidae

This S. American harvestman has spiny hind legs as a possible defence against predators. It lives under stones and logs in forests.

**HORNED HARVESTMAN***Phalangium opilio*

F: Phalangidae

A common harvestman of Eurasia and N. America, the male of this species has a jaw ornamented with projecting horns.

**TAILLESS WHIP SPIDER***Phrynus* sp.

F: Phrynidae

Whip spiders are not true spiders. They have long, whip-like front legs and pincers for grabbing prey, but lack venom. All are tropical.

**WHIP SCORPION***Thelyphonus* sp.

F: Thelyphonidae

Tropical whip scorpions such as this species have a whip-like tail, but no sting or venom. However, they can spray acetic acid from their abdomen.