


20–36 cm
8–14 in

TIGER PRAWN

Penaeus monodon
F: Penaeidae

This widely farmed, giant Indo-Pacific prawn belongs to a group that does not brood its fertilized eggs, but releases them directly into the sea.

PEACOCK MANTIS SHRIMP

Odontodactylus scyllarus
F: Odontodactylidae

Mantis shrimps of warm marine shallows are intelligent and formidable predators. This Indo-Pacific species uses “clubs” on its front limbs to smash the shells of crabs and snails.

3–18 cm
1½–7 in

carapace

stalked eyes

club-like
limbs deliver
powerful blows

3–5 cm
1¼–2 in

COMMON SHRIMP

Crangon crangon
F: Crangonidae

This shallow-water crustacean is commercially fished on some European coasts. It habitually hides in sand with only its eyes and antennae showing.


5–20 cm
2–8 in

SCULPTURED SLIPPER LOBSTER

Parribacus antarcticus
F: Scyllaridae

Relatives of spiny lobsters, clawless slipper lobsters have flattened, plate-like antennae. This nocturnal species lives on tropical reefs with sandy bottoms.

SINGAPORE BAMBOO SHRIMP

Atyopsis moluccensis
F: Atyidae

From a family of mostly freshwater river shrimps, this is a filter-feeder from S.E. Asia that eats particles caught on its fan-like front legs.



5–7.5 cm
2–3 in

7–11 cm
2¾–4¾ in

COMMON PRAWN

Palaemon serratus
F: Palaemonidae

A large, commercially important member of a family of shrimps with a jagged-edged carapace, this species inhabits northeast Atlantic coasts, frequenting the intertidal zone.


2.5 cm
1 in

SPOTTED CLEANER SHRIMP


Periclimenes yucatanicus
F: Palaemonidae

This Caribbean crustacean lives in sea anemones and – like many other reef-dwelling shrimps – cleans visiting fish of their dead skin and parasites.

COMMON MARBLE SHRIMP

Saron marmoratus
F: Hippolytidae

A member of the hump-back shrimp family, this Indo-Pacific reef species is a nocturnal predator of smaller animals. It turns red at night.


5 cm
2 in

STRIP-LEG SPINY LOBSTER

Panulirus femoristriga
F: Palinuridae

Unlike true lobsters, spiny lobsters, such as this Indo-Pacific species, lack claws and have a heavy, spiny exoskeleton and long, stout antennae.



20–30 cm
8–12 in

4–5 cm
1½–2 in

AESOP SHRIMP

Pandalus montagui
F: Pandalidae


Males of this cold-water, North Atlantic shrimp sometimes change into females after 13–16 months. Other individuals remain male or female throughout their lives.

8–12 cm
3¼–4¾ in


WHITE-CLAWED CRAYFISH

Austropotamobius pallipes
F: Astacidae

Crayfish are small, mostly nocturnal, freshwater relatives of true lobsters. This is a European stream species and the only crayfish native to Britain.


long stiff
antennae

20–24 cm
8–9½ in


long spiny claws

NORWAY LOBSTER


Nephrops norvegicus
F: Nephropidae

This North Atlantic crustacean tunnels in mud, often cohabiting with goby fish. It is a small-bodied true lobster, commercially marketed as langoustine or scampi.

SERRATED SLIPPER LOBSTER

Ibacus brevipes
F: Scyllaridae

Like other slipper lobsters, this Indo-Pacific crustacean can swim rapidly away from predators by flapping its broad tail fan.


5–12 cm
2–4¾ in