


## TRUE BUGS

Abundant and widespread in terrestrial and aquatic habitats, members of the order Hemiptera range from minute, wingless insects to giant water bugs capable of catching fish and frogs. The mouthparts are used for piercing and sucking up liquids such as plant sap, dissolved prey tissues, or blood. Many species are plant pests and some transmit disease.



8–10 mm  
3/16–3/8 in

**SPITTLE BUG**  
*Aphrophora alni*  
F: Aphrophoridae

Commonly occurring on a wide range of trees and shrubs across Europe, this froghopper can vary in appearance from pale to dark brown.



9–11 mm  
1/32–3/8 in

**FROGHOPPER**  
*Cecopis vulnerata*  
F: Cercopidae

The nymphs of this conspicuously coloured European species live communally underground in a protective, frothy mass and feed on plant root sap.



1.3–1.8 cm  
1/2–3/4 in

**LEAFHOPPER**  
*Ledra aurita*  
F: Cicadellidae


Mottled coloration helps this flat-bodied leafhopper from N. Europe to blend with the lichen-covered bark of oak trees in its habitat.



6–8 mm  
1/4–5/16 in

**CICADELLA VIRIDIS**  
F: Cicadellidae


Found feeding on grasses and sedges in wet, boggy, or marshy areas in Europe and Asia, this leafhopper can also be found near garden ponds.



1–2 mm  
1/32–1/16 in

**GLASSHOUSE WHITEFLY**  
*Trialeurodes vaporariorum*  
F: Aleyrodidae

This small moth-like bug is found in temperate regions worldwide and can be a serious pest of glasshouse crops.



3–5 mm  
1/8–3/16 in

**AMERICAN LUPIN APHID**  
*Macrosiphum albifrons*  
F: Aphididae

Aphids such as this U.S. species can quickly infest plants because females can produce many offspring without fertilization.

**INDIAN CICADA**  
*Angamiana aetherea*  
F: Cicadidae


This cicada is found in India. As with all cicadas, males produce loud songs both for courtship and to signal aggression.

35–40 mm  
1 1/2 in



**ASSAM CICADA**  
*Platyleura assamensis*  
F: Cicadidae


Known from N. India, this species is also found in Bhutan and parts of China where it prefers temperate, deciduous forests.



8 cm  
3 1/4 in

**PEANUT-HEADED BUG**  
*Fulgora laternaria*  
F: Fulgoridae

This bug is found in C. and S. America and the West Indies. The bulbous head was once thought to glow.



6–8 mm  
1/4–5/16 in

**HORSE CHESTNUT SCALE**  
*Pulvinaria regalis*  
F: Coccidae

Although commonly found on the bark of horse chestnut trees, this European scale insect will also attack a range of other deciduous species.



1–1.2 cm  
3/8–1/2 in

**THORN BUG**  
*Umbonia crassicornis*  
F: Membracidae

Found in C. and S. America, almost the whole body of this bug is concealed under the enlarged, thorn-like pronotum.



2–3 mm  
1/16–1/8 in

**ASH PLANT LOUSE**  
*Psyllopsis fraxini*  
F: Psyllidae


Found commonly on ash trees, the nymphs of this species cause the edges of leaves to form red, swollen galls when they feed.



3–4 mm  
1/8–5/32 in

**THISTLE LACE BUG**  
*Tingis cardui*  
F: Tingidae

This bug occurs in most of Europe and feeds on spear, musk, and marsh thistles. The body is covered in powdery wax.



3.2 cm  
1 1/4 in

**WART-HEADED BUG**  
*Phrictus quinquepartitus*  
F: Fulgoridae

Also known as the dragon-headed bug, this species can be found in Costa Rica, Panama, Colombia, and parts of Brazil.

distinctive marking across forewings

brightly coloured hindwings

elongated head

distinct, dark marginal area

pale area at base of hindwing