


**POWDER POST BEETLE**

Lyctus opaculus
F: Bostrichidae

This N. American beetle breeds in dry wood and reduces it to a fine powder.

SEXTON BEETLE

Nicrophorus investigator
F: Silphidae

This species is found all over the northern hemisphere in woodland and grassland. It buries the carcasses of small animals and the female lays an egg on it, providing food for the larva.

HERCULES BEETLE


Dynastes hercules
F: Scarabaeidae

The largest species of rhinoceros beetle (genus *Dynastes*), this beetle feeds on rotting fruits in rainforests in C. and S. America. Its larvae breed in decaying wood.

GOLIATH BEETLE

Goliathus cacicus
F: Scarabaeidae

This species, the world's heaviest insect, lives in equatorial Africa, where the adults feed on ripe fruits or tree sap.

**STAG BEETLE**

Lucanus cervus
F: Lucanidae

This impressive beetle lives in woodland in S. and C. Europe. Its larvae develop over 4–6 years, mainly in decaying oak stumps. Males use their enlarged, antler-like mandibles in contests over females.

NET-WINGED BEETLE

Platycis minutus
F: Lycidae

This small beetle is associated with decaying wood in mature and ancient woodland across Eurasia.

**PHANAEUS DEMON**

F: Scarabaeidae

Native to C. America, this very variable beetle breeds in the dung of large herbivorous animals in grassland and pasture.

**SAP BEETLE**

Glischrochilus hortensis
F: Nitidulidae

Often found feeding at fermenting sap flows or ripe fruits, this W. European member of the pollen beetle family is associated with decaying trees such as birch.

DEVIL'S COACH HORSE


Staphylinus olens
F: Staphylinidae

A European species found among woodland and garden leaf litter, this beetle raises its abdomen in a threatening display if disturbed.

**EMUS HIRTUS**

F: Staphylinidae

Native to S. and C. Europe, this hairy rove beetle feeds on other insects attracted to cow and horse manure or carrion.

**GOLD BEETLE**

Chrysina aurigans
F: Scarabaeidae

Found in Costa Rica and Panama, this beetle lives in deciduous forests, usually in rotting logs. The adults are nocturnal and attracted to light.

