

» BEETLES



3 mm
1/8 in

ANT-LIKE BEETLE
Anthicus floralis
F: Anthicidae

Found across Europe and parts of Asia, this small beetle lives in and around decaying vegetation, manure, and compost heaps.



1–1.2 cm
3/8–1/2 in

**THICK-LEGGED
FLOWER BEETLE**
Oedemera nobilis
F: Oedemeridae


Often seen feeding on pollen at flowers in meadows, this beetle is native to S.W. Europe. Its larvae develop inside plant stems.



2.4–3.4 cm
1–1 1/2 in

BLACK OIL BEETLE
Meloe proscarabaeus
F: Meloidae

This European beetle is found in warm meadows, heaths, and coastal areas. The larvae develop inside the nests of wild bees.




5–6 cm
2–2 1/4 in

**RED-SPOTTED
LONGHORN BEETLE**
Batocera rufomaculata
F: Cerambycidae

Also known as the mango borer or fig borer, this tropical species can be found throughout India and S.E. Asia.

1.8–2.2 cm
1/2–3/4 in



**GIBBIFER
CALIFORNICUS**
F: Erotylidae

This species is found in moist woodland in the S.W. USA. The larvae feed on fungi growing on trees and decaying wood.



1.4–1.6 cm
3/4 in

CARDINAL BEETLE
Pyrochroa coccinea
F: Pyrochroidae


This distinctive and somewhat flattened European beetle is often seen crawling on old tree stumps, inside which its larvae develop.



2.8–3.6 cm
1 1/4–1 1/2 in

**PHOSPHORUS
VIRESCENS**
F: Cerambycidae

Native to W. Africa, the larvae of this beetle burrow into some economically important species such as the cola tree.



4.2–5.5 cm
1 1/2–2 1/4 in

**WALLACE'S
LONGHORN BEETLE**
Cyriopalus wallacei
F: Cerambycidae

This impressive beetle can be found in the rainforests of S.E. Asia, where its larvae bore into certain trees.



3.2–5 cm
1 1/2–2 in


**DICRONORHINA
DERBYANA**
F: Cetoniidae

This beetle is found in southern and eastern parts of Africa, where the larvae feed on decaying plant matter and animal dung.

spines on
leg joint

TAURHINA POLYCHROUS
F: Cetoniidae

Native to E. Africa, especially Tanzania, this beetle exists in a number of colour forms. The blue form is considered to be a subspecies.



3–3.5 cm
1 1/4–1 1/2 in

large cephalic horn