



3–3.5 cm
1¼–1½ in

JEWELLED FROG BEETLE

Sagra buqueti
F: Chrysomelidae

This beetle develops inside certain large-stemmed vines in S.E. Asia, particularly Thailand. Like many attractive tropical species, they are sold to collectors.




6–8 mm
¼–⅝ in

SCARLET LILY BEETLE

Lilioceris lili
F: Chrysomelidae

Originally from Europe and Asia, this striking pest has spread to other parts of the world where lilies are grown.



7–9 mm
⅞–1½ in

EYED LADYBIRD

Anatis ocellata
F: Coccinellidae

This ladybird is native to Europe and is associated with coniferous trees, especially spruce and pine, where it feeds on aphids.



5–8 mm
⅜–⅝ in

SEVEN-SPOT LADYBIRD

Coccinella septempunctata
F: Coccinellidae

This species is very common in a wide range of habitats right across Europe, and is now established in N. America.




3–5 mm
⅛–⅜ in

BROAD BEAN WEEVIL

Bruchus rufimanus
F: Chrysomelidae

This species is a serious pest of broad bean crops. The adults feed on the pollen and their larvae tunnel into the seeds.



4–5 mm
⅝–⅜ in

LARCH LADYBIRD

Aphidecta obliterata
F: Coccinellidae

This European species lives on conifers such as larch, fir, and pine, where it feeds on scale insects and conifer aphids.



3–4 mm
⅛–⅝ in

TWENTY-TWO SPOT LADYBIRD

Psyllobora vigintiduopunctata
F: Coccinellidae

This European species lives on low-growing plants in meadows and, unlike most ladybirds, feeds on fungi such as mildews.



2–3 cm
¾–1¼ in



2.5–4 mm
⅛–⅝ in

BAPTISIA WEEVIL

Trichapion rostrum
F: Brentidae

This small weevil feeds on the seeds of white and cream wild indigo (*Baptisia*) plants on the prairies of N. America.



1–1.3 cm
⅜–½ in

CRATOSOMUS RODDAMI

F: Curculionidae

This species lives in C. and S. America. Its larvae are stem and bark borers of certain wild fruit trees.

GIRAFFE-NECKED WEEVIL

Trachelophorus giraffa
F: Attelabidae

This beetle lives in the rainforests of Madagascar. It rolls the leaves of the *Dichaetanthera* tree with its neck, and the female lays her egg inside the roll.

1.8–2 cm
⅞–¾ in



ONYMACRIS CANDIDIPENNIS

F: Tenebrionidae

Long legs and white wing cases help this diurnal beetle survive in the arid deserts along the southwestern coast of Africa.



2.6 cm
1 in



2–2.4 cm
¾–1 in

CHURCHYARD BEETLE

Blaps mucronata
F: Tenebrionidae

This flightless, ground-living species is usually active after dark and can be found in dark and damp places, where it eats decaying matter.

BLUE BANDED WEEVIL

Eupholus linnei
F: Curculionidae

Found on certain islands in E. Indonesia, this spectacular weevil, like many others in the genus *Eupholus*, feeds on yams.

8–12 mm
⅝–½ in



BLACK VINE WEEVIL

Otiorhynchus sulcatus
F: Curculionidae

Widespread in Europe, N. America, and Australasia, this weevil is a serious pest of a wide range of garden and cultivated plants.