


» TRUE FLIES

YELLOW DUNG FLY*Scathophaga stercoraria*
F: Scathophagidae

This very common fly is found in many parts of the northern hemisphere, where it breeds in cattle and horse dung. The larvae feed on the dung, while the adults prey on other insects attracted to the dung.

6–11 mm
¼–⅜ infleshy, toothed
proboscis for
attacking preyyellow, bristly
body1.3–1.5 cm
½–¾ in**BANDED BROWN HORSE FLY**
Tabanus bromius

F: Tabanidae


Widespread in Europe and the Middle East, this fly mainly attacks horses but will feed on some other animals, including humans.

3–5 mm
⅛–⅜ in**SEPSIS SP.**
F: Sepsidae


These flies are common and widespread in a wide variety of habitats. The larvae develop in animal dung and rotting matter.

2–3 mm
⅛–¼ in**COMMON FRUIT FLY**
Drosophila melanogaster
F: Drosophilidae


A common laboratory animal, this widespread species has a distinctive dark patch on the abdomen and breeds in rotting fruits.

8–14 mm
¼–¾ in**MARSH SNIPE FLY**
Rhagio tringarius
F: Rhagionidae


This predatory fly can be found on low vegetation growing on damp scrubby or boggy ground throughout much of Europe.

1–1.2 cm
⅜–½ in**SYRPHUS RIBESII**
F: Syrphidae


The larvae of this nectar-feeding fly are major predators of aphids in the northern hemisphere. Its coloration mimics that of wasps and bees to deter predators.

0.9–1.2 cm
⅜–½ in**MARMALADE HOVER FLY**
Eisyrphus balteatus
F: Syrphidae

Very common in a variety of habitats including gardens, this European fly feeds on pollen and nectar. The larvae eat aphids.

1–1.2 cm
⅜–½ in**LEUCOZONA LEUCORUM**
F: Syrphidae

This northern hemisphere species can be found visiting flowers in damp woodlands in the spring and early summer. Its larvae prey on aphids.

1.5–2 cm
⅝–¾ in**DRONE FLY**
Eristalis tenax
F: Syrphidae


This European fly, which has been introduced to N. America, is a good mimic of the honey bee. The larvae breed in stagnant water.

8 mm
⅜ in**FOREST FLY**
Hippobosca equina
F: Hippoboscidae

Occurring mainly in wooded areas in Europe and parts of Asia, this bloodsucking fly attacks horses, deer, and sometimes cattle.

4–5 mm
⅝–¾ in**PENICILLIDIA FULVIDA**
F: Nycteribiidae

Widespread in sub-Saharan Africa, this bloodsucking, wingless, ectoparasitic fly can be found on a wide range of bat species.

3.5–7 cm
1½–2¾ in**GAUROMYDAS HEROS**
F: Mydidae

This S. American insect, the world's largest fly, breeds inside the nests of leaf-cutter ants, where the young are thought to eat scarab beetle larvae.