




2.4–3 cm
1–1¼ in

RODENT BOT FLY
Cuterebra fontinella
F: Oestridae

This parasitic fly lives in N. America. The larvae develop inside the bodies of rodents such as white-footed mice.



4.5–5.5 mm
⅓–⅜ in

APPLE MAGGOT
Rhagoletis pomonella
F: Tephritidae


Native to N. America, this fly is an apple pest but also attacks hawthorn, plums, cherries, and other related fruit trees.



8–15 mm
¼–⅓ in

**SAVANNAH
TSETSE FLY**
Glossina morsitans
F: Glossinidae

Confined to parts of Africa, this fly is one of the species that transmits the trypanosome parasite that causes sleeping sickness in humans.



4–6 mm
⅝–¼ in

SNAIL-KILLING FLY
Pherbellia cinerella
F: Sciomyzidae

This species is found in damp meadows and marshy areas across Europe and parts of Asia. The larvae prey on snails of the family Helicidae.



3–5 cm
1¼–2 in

**PANTOPHTHALMUS
BELLARDII**
F: Pantophthalmidae

Found in the rainforests of Costa Rica, Venezuela, and Ecuador, the larvae of this large fly feed inside tree roots.



1–1.2 cm
⅜–½ in

DANCE FLY
Empis tessellata
F: Empididae

Native to Europe and Asia, this fly can be found in meadows and hedgerows, where it feeds on nectar and other insects.

CADDISFLIES

Closely related to the Lepidoptera, these slender, rather moth-like insects of the order Trichoptera are covered with hairs, not scales. The head has long, thread-like antennae and weakly developed mouthparts. The two pairs of wings are held tent-like over the body at rest. Caddisfly larvae are aquatic and often construct species-specific portable shelters out of small stones or plant fragments.



3–4.5 mm
⅓–⅓ in

SALT AND PEPPER MICROCADDIS
Agraylea multipunctata
F: Hydroptilidae

This small caddisfly is widespread and common in N. America. It has narrow wings and breeds in algae-rich ponds and lakes.



2.8–3.2 cm
1⅓–1¼ in

GREAT RED SEDGE
Phryganea grandis
F: Phryganeidae

This European caddisfly breeds in weedy lakes and slow rivers. The larvae make a case from cut leaf sections arranged in a spiral.



1.2–1.5 cm
½–⅓ in

MARBLED SEDGE
Hydropsyche contubernalis
F: Hydropsychidae


This caddisfly flies in the evening and breeds in rivers and streams. The aquatic larvae weave a net to catch food.



8–13 mm
¼–½ in

DARK-SPOTTED SEDGE
Philopotamus montanus
F: Philopotamidae

This European caddisfly has shortish antennae and breeds in fast-flowing rocky streams. The larvae make tube-like nets on the undersides of rocks.



1.2–1.7 cm
½–⅓ in

MOTTLED SEDGE
Glyptotendipes pellucidus
F: Limnephilidae

This European caddisfly breeds in lakes and small ponds. The larvae make a case out of fragments of withered tree leaves.