


## MOTHS AND BUTTERFLIES

Members of the order Lepidoptera are covered with minute scales. They have large compound eyes and a feeding tube called a proboscis. Larvae, known as caterpillars, pupate, metamorphosing into their adult form. Most moths are nocturnal and rest with their wings open; the majority of butterflies are diurnal and rest with their wings closed.

### HERCULES MOTH

*Coscinocera hercules*  
F: Saturniidae

Found in New Guinea and Australia, this moth is one of the biggest in the world. Only the male has the long hindwing tails.



20–27 cm  
8–10½ in

hindwing tail

10–16 cm  
4–6½ in

### AMERICAN MOON MOTH

*Actias luna*  
F: Saturniidae

This lime-green N. American moth has distinctive long hindwing tails. The caterpillars feed on a range of deciduous trees.



10–16 cm  
4–6½ in

### POLYPHEMUS MOTH

*Antheraea polyphemus*  
F: Saturniidae

Common and widespread in the USA and S. Canada, this moth has large eyespots on the wings, to startle predators.



6–9 cm  
2¼–3½ in

### GIANT LEOPARD MOTH

*Hypercompe scribonia*  
F: Erebidæ

The distribution of this striking moth extends from S.E. Canada as far south as Mexico. The caterpillars feed on a variety of plants.



4.5–7.5 cm  
1¾–3 in

### OAK EGGAR

*Lasiocampa quercus*  
F: Lasiocampidae

Ranging from Europe to N. Africa, the caterpillars of this moth feed on the foliage of bramble, oak, heather, and other plants.



5–8.5 cm  
2–3¼ in

### LAPPET MOTH

*Gastropacha quercifolia*  
F: Lasiocampidae

Found in Europe and Asia, the specific name of this large moth refers to its resting appearance, which resembles a cluster of dead oak leaves.



5–8 cm  
2–3¼ in

### PINE-TREE LAPPET

*Dendrolimus pini*  
F: Lasiocampidae

This moth is widespread in coniferous forests across Europe and Asia, where the caterpillars feed on pine, spruce, and fir.



3.5–4.5 cm  
1½–1¾ in

### AUSTRALIAN MAGPIE MOTH

*Nyctemera amicus*  
F: Erebidæ

Widespread in Australia and also found in New Zealand, the caterpillars of this day-flying moth feed on plants such as groundsel and ragwort.




4.5–7 cm  
1¾–2¾ in

### GARDEN TIGER

*Arctia caja*  
F: Erebidæ

This unmistakable moth is found across the northern hemisphere. The hairy caterpillars feed on a range of low-growing plants and shrubs.



7–8 cm  
2¾–3¼ in

### ILIA UNDERWING

*Catocala ilia*  
F: Erebidæ

This widespread species from N. America has distinctive red-banded hindwings. The caterpillars feed on the foliage of oak trees.




2.5–3 cm  
1–1¼ in

### VAPOURER MOTH

*Orgyia antiqua*  
F: Erebidæ

Now found across the northern hemisphere, the female of this European species has tiny wings and is unable to fly.



1–1.6 cm  
¾–1½ in

### CASE-BEARING CLOTHES MOTH

*Tinea pellionella*  
F: Tineidae

Found in W. Europe and parts of N. America, this species can cause serious damage to woollen clothing and carpets.