



24–30 cm
9½–12 in

GIANT AGRIPPA
Thysania agrippina
F: Noctuidae


Found in C. and parts of S. America, this species has one of the largest wingspans of any moth in the world.

SNOUT MOTH
Vitessa suradeva
F: Pyralidae

This moth occurs in India, parts of S.E. Asia, and New Guinea. The caterpillars feed in a web on young leaves of poisonous shrubs.



4–5 cm
1½–2 in



2.4–2.8 cm
1–1¼ in

SMALL MAGPIE
Anania hortulata
F: Pyralidae

Found in hedgerows and waste ground, the caterpillars of this very common European species feed on rolled nettle leaves.



6–10 cm
2¼–4 in

SILVER-SPOTTED GHOST MOTH
Sthenopis argenteomaculatus
F: Hepialidae

This species occurs in S. Canada and parts of the USA. The caterpillars mainly feed inside the roots of alder trees.

ACACIA CARPENTER MOTH
Endoxyla encalypti
F: Cossidae

The robust, white caterpillars of this large and distinctive Australian moth burrow into the woody tissue of certain species of the acacia tree.



9–12 cm
3½–4¾ in

DIVA MOTH
Divana diva
F: Castniidae

Found in tropical forests in S. America, this day-flying species is well camouflaged at rest despite its brightly coloured hindwings.



6.5–9.5 cm
2½–3¾ in

large eyespot to scare predators




5–6.5 cm
2–2½ in



LARGE EMERALD
Geometra papilionaria
F: Geometridae

This moth occurs across Europe and the temperate parts of Asia. The caterpillars feed mainly on the foliage of birch trees.


3–4 cm
1¼–1½ in



ARGENT AND SABLE
Rheumaptera hastata
F: Geometridae

The common name for this day-flying moth from the northern hemisphere comes from the heraldic terms for silvery-white and black.

4–5 cm
1½–2 in



CLARA'S SATIN MOTH
Thalaina clara
F: Geometridae


The caterpillars of this moth, which occurs in E. and S.E. Australia and N. Tasmania, feed on acacia foliage.

wavy pattern of black and orange lines

10–16 cm
4–6½ in

WALLICH'S OWL MOTH
Brahmaea wallichii
F: Brahmaeidae

This large moth occurs from N. India to China and Japan. The caterpillars eat the foliage of ash, privet, and lilac.




7–7.5 cm
2¾–3 in

COPPERY DYSPHANIA
Dysphania cuprina
F: Geometridae

Widespread across S.E. Asia, this brightly coloured moth flies during the day and is thought to be unpalatable to birds.

SILK MOTH
Bombyx mori
F: Bombycidae


Originating in China, the silk moth has been bred in captivity, on a diet of white mulberry leaves, for thousands of years. Its cocoon is used to make silk.



4–6 cm
1½–2¼ in

WHITE PLUME-MOTH
Pterophorus pentadactyla
F: Pterophoridae

Common across Europe on dry grassland, waste ground, and gardens, the caterpillars of this distinctive species feed on hedge bindweed.



2.5–3 cm
1–1¼ in

long legs