




8–14 cm
3¼–5½ in


TIGER SWALLOWTAIL
Papilio glaucus
F: Papilionidae

This species is widespread in N. America. The young caterpillars, which resemble bird droppings, feed on a variety of trees and shrubs.



SMALL COPPER
Lycaena phlaeas
F: Lycaenidae

Common in Europe, N. Africa, and Asia as far as Japan, this species can also be found in N. America. The caterpillars feed on sorrels and docks.



2.5–3 cm
1–1¼ in

BROWN HAIRSTREAK
Thecla betulae
F: Lycaenidae

This species can be found in hedgerows, scrubland, and woodland across Europe and temperate Asia. The caterpillars feed at night on blackthorn.



3–4 cm
1¼–1½ in

BLUE THAROPS
Menander menander
F: Lycaenidae

This fast-flying butterfly occurs in tropical forests from Panama to northern S. America. Little is known about its life cycle or caterpillars.



DUKE OF BURGUNDY FRITILLARY
Hamearis lucina
F: Lycaenidae

This species ranges across C. Europe as far as the Urals and favours flower meadows, where its food plants, cowslip and primrose, grow.



2.5–3.5 cm
1–1½ in

ADONIS BLUE
Lyandra bellargus
F: Lycaenidae

This European species can be found in chalky grassland, where the caterpillars feed on horseshoe vetch. The male is sky blue while the female is brown.



2–2.5 cm
¾–1 in

SONORAN BLUE
Philotes sonorensis
F: Lycaenidae

This rare butterfly is restricted to rocky washes and desert cliffs in California. The caterpillars feed on succulents known as stonecrop or “live-forever” plants.

SAWFLIES, WASPS, BEES, AND ANTS

Hymenoptera typically have two pairs of wings, joined in flight by tiny hooks. Apart from sawflies, they have a “wasp-waist” and females have an ovipositor that may be modified to sting. Many wasps are predatory or parasitic. Bees are vital pollinators and ants play a role in many ecosystems.



7–9 mm
7/32–1½/32 in

fine tufts on wing margin



TENTHREDO ARCUATA
F: Tenthredinidae

Widespread in meadows, this distinctive black- and yellow-marked European sawfly lays its eggs on clovers.



1.8–2.2 cm
¾–9/10 in

CIMBICID SAWFLY
Trichiosoma lucorum
F: Cimbicidae

This stout-bodied European sawfly is found in woodland, hedgerows, and scrubby areas. The larvae feed on birch and willow.



7–9 mm
7/32–1½/32 in

STEM SAWFLY
Cephus nigrinus
F: Cephidae

This slender, all-black sawfly is widespread in W. Europe. The larvae bore downwards inside the stems of wild and some cultivated grasses.



2 cm
¾ in

ROSE SAWFLY
Arge ochropus
F: Argidae

Recognizable by the black along the wing edge, this European sawfly visits flat-topped flowers to feed. The larvae feed on wild roses.



7–10 mm
7/32–¾ in

PERGID SAWFLY
Unknown sp.
F: Pergidae


These herbivorous insects are found in Australia and S. America. Many species attack eucalyptus trees, and young larvae feed gregariously.



7–9 mm
7/32–1½/32 in

RED-HEADED SAWFLY
Acantholyda erythrocephala
F: Pamphiliidae

Originally found in Europe and Asia, this sawfly has spread to North America. The larvae feed communally under silken webs among pine needles.



HORNTAIL
Urocerus gigas
F: Siricidae

This impressive species can be found across the northern hemisphere. The female drills deep into softwood trees to lay her eggs.

ovipositor