

» PERIWINKLES, WHELKS, AND RELATIVES



10–32 cm
4–12½ in

GIANT FROG SHELL
Tutufa bubo
F: Bursidae

Tropical marine frog shells are warty. This Indo-Pacific species eats bristleworms using a proboscis, first anaesthetizing them with saliva.

warty shell

canal for siphon



7.5–11.5 cm
3–4½ in

BANDED TULIP
Cinctura lilium
F: Fascioliariidae


This is a Caribbean snail related to *Buccinum* whelks. Other sand-dwelling species with a longer siphonal canal are called spindle shells.



6–11 cm
2¼–4¼ in

IMPERIAL HARP
Harpa costata
F: Harpidae

Harp snails are sand-living predators of crabs, trapping them with their broad foot and digesting them with saliva. This is an Indian Ocean species.



2–3 cm
¾–1¼ in

PRICKLY PACIFIC DRUPE
Drupa ricinus
F: Muricidae

A member of the dog whelk family, this snail lives on coral reefs of the Indo-Pacific, where it preys on bristle worms.



3–4 cm
1¼–1½ in

DOG WHELK
Nucella lapillus
F: Muricidae

This North Atlantic snail belongs to a whelk family that catch prey by using the radula to drill holes in the shells of barnacles and molluscs, helped by secretions of enzymes.



9–15 cm
3½–6 in

TEXTILE CONE
Conus textile
F: Conidae

Cone shells use their radula to harpoon their prey and inject venom. Some, such as this Indo-Pacific species, can be harmful to humans.



3–13 cm
1½–5 in

TENT OLIVE
Oliva porphyria
F: Olividae

The largest of the olive shells, this species from the Pacific coasts of Mexico and S. America has a colourful glossy shell.



10–50 cm
4–20 in

TRUMPET TRITON
Charonia tritonis
F: Charoniidae

Tropical intertidal tritons and related frog and helmet shells are predators of other invertebrates. This Indo-Pacific species eats the coral-feeding crown-of-thorns starfish.



5 mm–3 cm
¾–1¼ in

POLI'S NECKLACE SHELL
Euspira nitida
F: Naticidae

This is a European member of a family of sand-burrowing predators of bivalves. Necklace shells are named for their necklace-like ribbons of spawn.



8–11 cm
3¼–4¼ in

COMMON NORTHERN WHELK
Buccinum undatum
F: Buccinidae

A large predator of other molluscs and tube worms, this North Atlantic snail also consumes carrion, and is a popular seafood.




7–20 cm
2¼–8 in

SUBULATE AUGER
Terebra subulata
F: Terebridae

Typical of many augers, this Indo-Pacific species has a patterned shell. Augers burrow through surface layers of sand to prey on worms.

FRESHWATER GILLED SNAILS

The Architaenioglossa are the only order of gilled snails containing no marine species. Most live in fresh water, but some live on land. Apple snails have gills in the mantle cavity that can function as a lung, allowing them to survive periods of drought. All species have an operculum, or trap door, for closing their shell.



10–15 cm
4–6 in

CHANNELLED APPLE SNAIL
Pomacea canaliculata
F: Ampullariidae

A typical apple snail, this tropical American species has been introduced elsewhere and has become an invasive pest.

muscular foot

COMMON RIVER SNAIL
Viviparus viviparus
F: Viviparidae

A European freshwater snail with gills, this is a relative of the apple snails and, like them, has a trapdoor (operculum) to close its shell.

3–4 cm
1¼–1½ in

