

flattened head
and body

1.5 m
5 ft

SAND DEVIL
Squatina dumeril
F: Squatinidae

Sandy seabeds in the N.W. Atlantic are home to this well-camouflaged predator. It strikes at passing fish with lightning speed.


gill
slit

ANGELSHARKS

Flattened from top to bottom, angelsharks have gill slits on the sides of their large head. This distinguishes them from the similarly shaped rays, which have gills underneath. The order Squatiniformes contains just one family, which has 25 members. Angelsharks rear up from the seabed on their pectoral fins to ambush prey.

GROUND SHARKS


There are about 295 species of Carcharhiniformes, making this the largest and most diverse order of sharks. Most are large predators, although the small yet numerous cat sharks are also included in the order. All have two dorsal fins and one anal fin.



4 m
13 ft

BLUE SHARK
Prionace glauca
F: Carcharhinidae


Streamlined and elegant, this species travels vast distances between feeding and breeding grounds. More blue sharks are caught than any other shark.



2 m
6½ ft

WHITETIP REEF SHARK
Triaenodon obesus
F: Carcharhinidae


This is perhaps the most common shark on the coral reefs of the Indian and Pacific oceans. It hunts fish at night, sometimes in frenzied packs.



4 m
13 ft

OCEANIC WHITETIP SHARK
Carcharhinus longimanus
F: Carcharhinidae

Heavily fished for its long, white-tipped dorsal and pectoral fins, this magnificent oceanic predator is now critically endangered.




3.4 m
11 ft

TIGER SHARK
Galeocerdo cuvier
F: Carcharhinidae

Named for its stripes, which fade with age, the tiger shark will eat anything and has attacked humans. Its serrated teeth are shaped like a cockscomb.

BULL SHARK
Carcharhinus leucas
F: Carcharhinidae

One of the ocean's most dangerous sharks, this top predator inhabits tropical coastal waters worldwide and also swims up rivers.



7.5 m
25 ft

rear
dorsal fin

vertical dark
bars and spots

anal fin

broad blunt
snout

pelvic fin