


ANGLERFISH AND RELATIVES

The 376 or so Lophiiformes include some of the most bizarre of all marine fish. A modified fin ray on top of the head acts as a fishing lure (bioluminescent in deep-water species), which attracts prey towards the fish's cavernous mouth.


COFFINFISH
Chaunax endeavouri
F: Chaunacidae
Coffinfish lie on muddy seabeds in the S.W. Pacific, waiting for small fish to stray within reach.



RED-LIPPED BATFISH
Ogcocephalus darwini
F: Ogcocephalidae
The oddly shaped batfish scrambles over the seabed using its paired pectoral and pelvic fins as props. Its red lips are a mystery.

ANGLER
Lophius piscatorius
F: Lophiidae

A fringe of seaweed-shaped flaps around its mouth helps disguise this N.E. Atlantic angler. It can strike with lightning speed.




WARTY FROGFISH
Antennarius maculatus
F: Antennariidae

The well-camouflaged warty frogfish clambers over coral reefs using its limb-like pectoral fins.


FANFIN ANGLER
Caulophryne jordani
F: Caulophrynidae

In the dark ocean depths it is difficult to find a mate. Once successful, the tiny male fanfin angler latches on to the female permanently.



SARGASSUMFISH
Histrion histrio
F: Antennariidae

Whilst most frogfish live on the seabed, this one hides amongst rafts of floating *Sargassum* seaweed, which it closely resembles.



large pectoral fins used for scrambling

skin flaps for camouflage


COD AND RELATIVES

The Gadiformes include many important marine commercial species. Most have two or three soft dorsal fins on their back and many have a chin barbel. Grenadiers live in deep water and have a long thin tail.



ATLANTIC COD
Gadus morhua
F: Gadidae

Overfishing has reduced the average weight of Atlantic cod to 11 kg (24 lb) from a historical maximum weight of over 90 kg (200 lb).



ALASKA POLLOCK
Gadus chalcogrammus
F: Gadidae

The lack of a chin barbel and a protruding lower jaw help distinguish this fish from cod. It lives in cold Arctic waters.



SHORE ROCKLING
Gaidropsarus mediterraneus
F: Lotidae

Equipped with three sensory mouth barbels, this eel-like rockling searches out food in N.E. Atlantic rock pools.



BURBOT
Lota lota
F: Lotidae

Unlike all other Gadiformes, the burbot lives in fresh water. It is found in deep lakes and rivers across the northern hemisphere.



PACIFIC GRENADIER
Coryphaenoides acrolepis
F: Macrouridae

A long scaly tail and bulbous head give this abundant deep-water cod relative its alternative name of Pacific rat-tail.