

## » PERCH AND RELATIVES

1.5 m  
5 ft

**WOLF-FISH**  
*Anarhichas lupus*  
F: Anarhichadidae

A large mouth and strong teeth enable this North Atlantic species to tackle hard-shelled invertebrates.

15 cm  
6 in

**LESSER WEEVERFISH**  
*Echiichthys vipera*  
F: Trachinidae

This venomous N.E. Atlantic fish lies buried in the sand and can inflict a painful sting from its spiny, black, first dorsal fin.

**SMALL SANDEEL**  
*Ammodytes tobianus*  
F: Ammodytidae

This and other small sandeel species are vital food source for seabirds. It is found in large shoals in shallow sandy bays of the N.E. Atlantic Ocean.

20 cm  
8 in

**CUCKOO WRASSE**  
*Labrus mixtus*  
F: Labridae

Blue-marked males of the species (shown here) develop from large, pink females with black and white tail markings, who change both colour and sex. This fish is found in the N.E. Atlantic and Mediterranean.




**BLACK SWALLOWER**  
*Chiasmodon niger*  
F: Chiasmodontidae

Found in deep tropical and subtropical waters, where food is scarce, this fish, with its huge mouth and distensible stomach, can tackle prey larger than itself.

25 cm  
10 in

**CLEANER WRASSE**  
*Labroides dimidiatus*  
F: Labridae

This tiny wrasse forms a mutualistic relationship with other reef fish. It cleans and eats parasites off these fish at specially set up "cleaner stations" on the reef.

14 cm  
5½ in


**ANTARCTIC TOOTHFISH**  
*Dissostichus mawsoni*  
F: Nototheniidae

Like many Southern Ocean fish, this species grows slowly, eventually reaching a large size. Its blood contains natural antifreeze, which helps it to survive icy waters.

2.2 m  
7¼ ft

**MAORI COD**  
*Paranotothenia magellanica*  
F: Nototheniidae

Found as far north as New Zealand and Argentina, this Southern Ocean species is commercially important as food.

38 cm  
15 in

**DOLPHINFISH**  
*Coryphaena hippurus*  
F: Coryphaenidae

This fast predator of the open ocean can reach a speed of 60 kph (37 mph). It is found worldwide in warm waters.

2.1 m  
7 ft

**LONGNOSE HAWKFISH**  
*Oxycirrhites typus*  
F: Cirrhitidae

This Indo-Pacific species stays well camouflaged among sea fans and black corals, where it preys on small invertebrates.

13 cm  
5 insmall mouth  
for picking out  
invertebratespattern  
provides  
camouflage1.2 m  
4 ft


**BIGEYE TREVALLY**  
*Caranx sexfasciatus*  
F: Carangidae

Built for speed, this predator hunts other fish at night over reefs in the Indian and Pacific oceans. Juveniles live inshore and may enter estuaries.



**SHARKSUCKER**  
*Echeneis naucrates*  
F: Echeneidae

This species has a ridged sucker on top of its head that allows it to cling onto large fish, dolphins, and turtles. It benefits from scraps as its host feeds.

1 m  
3¼ ft

reddish coloration overall