

GLASS FROGS

Found in Central and South America, frogs of the family Centrolenidae are known as “glass frogs” because many species have transparent skin on the underside through which their internal organs are visible.




FLEISCHMANN'S GLASS FROG
Hyalinobatrachium fleischmanni

Males of this species are territorial, using calls to defend their territory and to attract females. The females lay eggs on leaves over water.

2–3 cm
¾–1¼ in

adhesive pad
at tip of digits



WHITE-SPOTTED COCHRAN FROG
Sachatamia albomaculata

This species is found in wet lowland forests, and breeds near streams. Males call to females from low vegetation nearby.

2–3 cm
¾–1¼ in



silver eyes with
black reticulations

LIMON GIANT GLASS FROG
Sachatamia ilex

This arboreal frog lives in wet vegetation near streams. It has dark green bones, visible through its skin.

2.5–3.5 cm
1–1½ in



2–3 cm
¾–1¼ in

EMERALD GLASS FROG
Espadarana prosoblepon

Males of this arboreal frog are fiercely territorial, defending their space by calling, and occasionally fighting rivals while hanging upside down.

CERATOPHRYIDAE

These South American horned frogs have a very large head and wide mouth that enable them to eat animals nearly as large as themselves. They are “sit-and-wait” predators, remaining still and well-camouflaged before their prey comes within range.

“horn” above eye

ORNATE HORNED FROG
Ceratophrys ornata

A voracious predator living in grasslands in Argentina, this species breeds after heavy rain, in temporary pools and ditches.



8–13 cm
3¼–5 in

CRANWELL'S HORNED FROG
Ceratophrys cranwelli

Spending much of its life underground, this large frog emerges after heavy rain to mate and lay its eggs in pools.

9–14 cm
3½–5½ in

4–10 cm
1½–4 in



BUDGETT'S FROG
Lepidobatrachus laevis

This frog has a flattened body, a wide mouth, and fangs. It spends dry periods in a cocoon below ground, emerging to breed after rain.

wide
mouth



ROBBER FROGS

Found in North, Central, and South America, the Craugastoridae produce eggs that develop directly into small adults, without a tadpole stage. Eggs may be deposited on the ground or in vegetation. Many species show parental care of the eggs.

BROAD-HEADED RAIN FROG
Craugastor megacephalus

This C. American frog hides in a burrow by day and emerges at night. It lays its eggs in leaf litter.

3–7 cm
1¼–2¾ in



ISLA BONITA ROBBER FROG
Craugastor crassidigitus

Native to the humid forests of C. America, this ground-living frog is also found in coffee plantations and pasture.



2–5 cm
¾–2 in



2.5–5.5 cm
1–2¼ in

COMMON RAIN FROG
Craugastor fitzingeri

In this forest species, the male calls to the larger female from an elevated perch. Eggs are laid in the ground and are guarded by the female.