




5–9 cm
2–3½ in

AMERICAN TOAD
Anaxyrus americanus

Found in eastern N. America, this toad is quite variable in colour. Breeding occurs in ponds where males produce long trilling calls.

large
parotoid
glands

short,
warty legs

TRUANDO TOAD
Rhaebo haematiticus
Living among leaf litter in forests of C. and S. America, this broad-headed toad lays its eggs in long strings in rocky pools.



4–8 cm
1½–3¼ in



8–20 cm
3¼–8 in

EUROPEAN COMMON TOAD
Bufo bufo

Found throughout Europe and N. Africa, this species breeds in spring, with males outnumbering the much larger females by about three to one.



5.5–9.5 cm
2¼–3¾ in

GREEN CLIMBING FROG
Incilius coniferus
Native to C. and S. America, this nocturnal toad is often found climbing among vegetation.



VARIABLE HARLEQUIN FROG
Atelopus varius

This aggressive species from Panama and Costa Rica has vivid and variable coloration. It lives near streams and is active during the day.


2.5–6 cm
1–2¼ in



2.5–4 cm
1–1½ in

GUYANAN STUBFOOT TOAD
Atelopus barbotini

This small toad from Guyana has a flattened body. It breeds throughout the year in forest streams.



5–10 cm
2–4 in

PANAMANIAN GOLDEN FROG
Atelopus zeteki

This brightly coloured frog from Panama breeds in pools after heavy rain. It may now be extinct in the wild.



parotoid glands
secrete toxin


olive-
brown,
warty skin

10–24 cm
4–9½ in

CANE TOAD
Rhinella marina
One of the world's largest toads, this American species was introduced in Australia, where it is now a serious threat to native wildlife.

RHINODERMATIDAE

This South American family includes two species with a pointed snout and cryptic coloration, and the males carry the eggs and tadpoles in their mouth. The remaining species is rare and little is known about it.



green
back

fleshy
proboscis

eyes with
horizontal
pupils


2–3 cm
¾–1¼ in

DARWIN'S FROG
Rhinoderma darwini

Found in Chile and Argentina, this frog has a unique form of parental care – the young develop from egg to froglet in their father's vocal sacs.


ELEUTHERODACTYLIDAE

This group of frogs, whose eggs develop directly into froglets, is found throughout the Caribbean, the southern USA, and northern South America. Some species that are extremely small have reduced numbers of digits, and produce very small numbers of eggs – sometimes only one.



1.5–8 cm
½–3¼ in

PUERTO RICAN COQUI
Eleutherodactylus coqui
Native to Puerto Rico, this frog gets its name from its two-part call – the “co” is a warning to other males and the “qui” attracts females.



1.5–2.5 cm
½–1 in

large toe
pads

CARETTA ROBBER FROG
Diasporus diastema

This small and very agile frog is arboreal and active at night. It lays its eggs in water-filled cavities in bromeliads growing epiphytically in trees.