




SOUTHERN CASSOWARY

Casuarus casuarus
F: Casuariidae

Cassowaries are fruit-eating rainforest birds. With a range that extends through New Guinea and N. Australia, this is the most widespread species of its family.

1.5–1.8 m
5–6 ft




EMU

Dromaius novaehollandiae
F: Dromaiidae

Australia's largest living bird is found across the continent in open grassy areas. Like the related cassowary, its chicks are striped for camouflage.

1.7–2.1 m
5½–7 ft



LESSER RHEA

Rhea pennata
F: Rheidae

Smaller than the greater rhea, its northern cousin, this ratite inhabits the S. Andes and Patagonia, where it lives in small flocks.

92–100 cm
3–3¼ ft



black body plumage

almost bare neck

white primary feathers

1.7–2.7 m
5½–8¾ ft

OSTRICH
Struthio camelus
F: Struthionidae

The largest birds in the world, ostriches live in savanna and semi-desert areas of Africa. Breeding males mate with several females.

large eye

scaly legs

featherless thighs

two-toed feet



GREATER RHEA

Rhea americana
F: Rheidae

This central S. American ratite has a breeding system that does not involve pair-bonding. Males tend a large nest of eggs produced by many females.

1.3–1.4 m
4¼–4½ ft




SOMALI OSTRICH

Struthio molybdophanes
F: Struthionidae

Separated from other populations of ostrich by the Rift Valley, this E. African grey-necked form is now regarded as a separate species.

grey neck

1.7–2.7 m
5½–8¾ ft




GREAT SPOTTED KIWI

Apteryx haastii
F: Apterygidae

One of two species of kiwis with grey mottled plumage, this bird is confined to mountain areas of western South Island.

65–70 cm
26–28 in



LITTLE SPOTTED KIWI

Apteryx owenii
F: Apterygidae

Introduced mammals have driven this small kiwi to the edge of extinction. It survives on predator-free satellite islands around New Zealand.

35–45 cm
14–18 in