

grey-brown
upperparts

48–53 cm
19–21 in

olive-brown
breast

46 cm
18 in

**PLAIN
CHACHALACA**
Ortalis vetula
F: Cracidae

Ranging from Texas to Costa Rica, this northernmost member of the American curassow family is the only one found in the USA.

**GREY-HEADED
CHACHALACA**
Ortalis cinereiceps
F: Cracidae

Named after their calls, chachalacas are brown-coloured relatives of curassows. This is a species of scrubby habitat from Honduras to Colombia.

**BLUE-THROATED
PIPING GUAN**
Pipile cumanensis
F: Cracidae

This S. American bird has glossy black plumage typical of tree-dwelling piping guans. They are named for their shrill breeding call.

69 cm
27 in

59–65 cm
23–26 in

HIGHLAND GUAN
Penelopina nigra
F: Cracidae

This C. American bird is more ground dwelling than other guans. It is perhaps the only member of its family to nest on the ground.

white-tipped tail

SPIX'S GUAN
Penelope jacquacu
F: Cracidae

This S. American species belongs to a group of brown guans that resemble chachalacas, but have shorter legs. It nests in trees and has a loud mating call.

76 cm
30 in

CAUCA GUAN
Penelope perspicax
F: Cracidae

A bronze-sheened relative of the spix's guan, this bird is found only in the Cauca Valley region in the western and northern parts of Colombia.

61–71 cm
24–28 in

bare
bluish head

BEARDED GUAN
Penelope barbata
F: Cracidae

Named for its streaked neck, the bearded guan, like related species in this genus, is a rainforest bird. It is found in Ecuador and Peru.

55 cm
22 in

67–75 cm
26–30 in

red throat
wattle

**DUSKY-LEGGED
GUAN**
Penelope obscura
F: Cracidae

The only brown guan with dark rather than reddish legs, this species occurs in central S. America, from Brazil to northern Argentina.

68–75 cm
27–30 in

**CHESTNUT-BELLIED
GUAN**
Penelope ochrogaster
F: Cracidae

The throat wattle that is typical of brown guans is especially well developed in this species from south-central Brazil.

cape of
long feathers

**VULTURINE
GUINEAFOWL**
Acryllium vulturinum
F: Numididae

Guineafowl are African gamebirds that live in groups, but are monogamous. They all have a bare head. This is the largest species of guineafowl and is found in E. Africa.

MOUNTAIN QUAIL
Oreortyx pictus
F: Odontophoridae

This native of the Rocky Mountains, like other American quails, is a ground-nesting, monogamous species.

26–31 cm
10–12 in

CALIFORNIAN QUAIL
Callipepla californica
F: Odontophoridae

Found from Oregon to California, this species is characterized by its forward-drooping crest, comprised of a cluster of six feathers.

24–27 cm
9½–10½ in

GAMBEL'S QUAIL
Callipepla gambelii
F: Odontophoridae

This relative of the Californian quail has a longer crest. It occurs in the desert in the south of California and the ranges of the two do not overlap.

25 cm
10 in