


» FOWL, GAMEBIRDS,
AND RELATIVES32–35 cm
12½–14 in**CHUKAR PARTRIDGE***Alectoris chukar*
F: Phasianidae

With a range extending across central Eurasia, this is the most widespread of the red-legged partridges – typically birds of dry country, with striking black markings.

29–32 cm
11½–12½ in**GREY PARTRIDGE***Perdix perdix*
F: Phasianidae


This is the most widespread of a small genus of dark-bellied Eurasian gamebirds that are more related to pheasants than other partridges.

16–18 cm
6½–7 in**COMMON QUAIL***Coturnix coturnix*
F: Phasianidae


Unusually for a gamebird, this tiny species of W. Eurasian grasslands and semi-desert is migratory in the northern parts of its range.

43 cm
17 in**GREATER PRAIRIE CHICKEN***Typanuchus cupido*
F: Phasianidae

This species is from the C. USA. As with other species of prairie chicken, males engage in group courtship displays by inflating their coloured neck sac.

28–30 cm
11–12 in**GREY FRANCOLIN***Francolinus pondicerianus*
F: Phasianidae

As with related junglefowl, males of this S. Asian francolin have leg spurs that they use in fighting.

25–38 cm
10–15 in**RED-NECKED FRANCOLIN***Francolinus afer*
F: Phasianidae

A ground-nesting bird of forest and grassland, this species is widespread through Africa, from the Democratic Republic of Congo to the Cape.

39–40 cm
15½–16 in**SPRUCE GROUSE***Canachites canadensis*
F: Phasianidae

Like other woodland grouse, this N. American species can digest the needles of conifer and spruce trees, rejected by most animals.

38–41 cm
15–16 in**LESSER PRAIRIE CHICKEN***Typanuchus pallidicinctus*
F: Phasianidae

Found in southern N. America, this small species of prairie chicken has a coloured neck sac and drumming call, typical of *Typanuchus* grouse.

38 cm
15 in**ARABIAN PARTRIDGE***Alectoris melanocephala*
F: Phasianidae

This large red-legged partridge occurs in semi-desert regions of the Arabian Peninsula and Yemen. Its preferred habitat is juniper woodland.

21 cm
8½ in**CHESTNUT-BELLIED HILL PARTRIDGE***Arborophila javanica*
F: Phasianidae


Found in S.E. Asian rainforests, hill partridges are small, cryptically coloured gamebirds with stumpy tails. Many, such as this Javan species, have distinctive head markings.

26 cm
10 in**ROULROUL***Rollulus rouloul*
F: Phasianidae


Related to hill partridges, the S.E. Asian rouleoul has a more colourful plumage. Males are dark blue and have a reddish crest; females are green and lack the crest.

31 cm
12 in**CHINESE BAMBOO PARTRIDGE***Bambusicola thoracicus*
F: Phasianidae

There are two species of bamboo partridge, birds of E. Asia that are related to francolins and junglefowl. This bird is native to China.

40–50 cm
16–20 in**SOOTY GROUSE***Dendragapus fuliginosus*
F: Phasianidae

One of several N. American grouse that have an inflatable neck sac, this dark-coloured species inhabits the pine forests of the Pacific coast.

41–47 cm
16–18½ in**SHARP-TAILED GROUSE***Typanuchus phasianellus*
F: Phasianidae

This N. American grouse is a more widespread, more northerly cousin of prairie chickens. Males have purple sacs for display.

