



46–55 cm
18–22 in



SURF SCOTER
Melanitta perspicillata
F: Anatidae

The N. American surf scoter, like other scoter species, breeds near fresh water and winters at sea. The male's body is entirely black.

42–50 cm
16½–20 in



HOODED MERGANSER
Lophodytes cucullatus
F: Anatidae

Found in N. America, this bird has a saw-edged bill for catching fish. It dives with powerful kicks of the feet.


46–54 cm
18–22 in



BRONZE-WINGED DUCK
Specularia specularis
F: Anatidae

This duck is found along the rivers of S. America. It is locally called "dog-duck" after the barking call of the female.

36–45 cm
14–18 in



PINK-EARED DUCK
Malacorhynchus membranaceus
F: Anatidae

The pink head spot of this widespread Australian species is less distinctive than its zebra pattern. It consumes plankton, strained from water by beak flaps.

35–43 cm
14–17 in



RUDDY DUCK
Oxyura jamaicensis
F: Anatidae

This N. American duck, now introduced in Europe, has a stiff tail which it uses as a rudder when diving.

40–47 cm
16–18½ in



TUFTED DUCK
Aythya fuligula
F: Anatidae

A Eurasian member of the pochard group, the tufted duck feeds mostly, though not entirely, on invertebrates in contrast to its largely vegetarian relatives.


48–61 cm
19–24 in



CANVASBACK
Aythya valisineria
F: Anatidae

This N. American bird is the largest species of pochard, a group of ducks with typically stocky bodies and large heads.


38–58 cm
15–23 in



LONG-TAILED DUCK
Clangula hyemalis
F: Anatidae

Unlike most other Arctic sea ducks, the long-tailed duck breeds in saltwater as well as freshwater habitats. Males have a distinctive long tail.

35–44 cm
14–17½ in



SMEW
Mergellus albellus
F: Anatidae

A member of the group of cavity-nesting mergansers, this species is the only small white duck found in northern Eurasia.

61–63 cm
24–25 in



COMMON SHELDUCK
Tadorna tadorna
F: Anatidae

This goose-like duck is largely a coastal resident in Europe, but in Asia it migrates southwards from inland regions during winter. It nests in burrows.

52–58 cm
20½–23 in



RED-BREASTED MERGANSER
Mergus serrator
F: Anatidae

Widespread across the northern hemisphere, the red-breasted merganser breeds on coasts and spends more time at sea than other mergansers.

43–63 cm
17–25 in



KING EIDER
Somateria spectabilis
F: Anatidae

This bird breeds along Arctic tundra coastlines. Its large body size may help it to dive deeply for invertebrate prey.


55–56 cm
22 in



ROSYBILL
Netta peposaca
F: Anatidae

This S. American duck is related to the diving pochards, but spends more time feeding at the water surface. Only the males have a red bill.

51–61 cm
20–24 in



CRESTED DUCK
Lophonetta specularioides
F: Anatidae

This Andean species is a possible relic of a S. American lineage that was ancestral to more widespread dabbling ducks, such as the mallard.

43–48 cm
17–19 in



STELLER'S EIDER
Polysticta stelleri
F: Anatidae

Like other related sea ducks of Arctic and subarctic regions, this eider overwinters further south in huge flocks, sometimes numbering 20,000 birds.

