

» ALBATROSSES, PETRELS, AND SHEARWATERS



30 cm
12 in

AUDUBON'S SHEARWATER
Puffinus lherminieri
F: Procellariidae

This small bird breeds on tropical oceanic islands. Different populations are considered by some to be separate species.



45–56 cm
18–22 in

CORY'S SHEARWATER
Calonectris diomedea
F: Procellariidae

A large shearwater that glides with bowed wings, this species breeds on islands in the eastern Atlantic and disperses more widely in winter.



43 cm
17 in

ANTARCTIC PETREL
Thalassoica antarctica
F: Procellariidae

A bird of sub-Antarctic waters, this large species dives for fish and squid. It breeds on islands around the Antarctic.



31 cm
12 in

JOUANIN'S PETREL
Bulweria fallax
F: Procellariidae

A tropical petrel of the northwestern Indian Ocean, this bird has a weaving flight. It is closely related to the gliding shearwaters.



48 cm
19 in

BULLER'S SHEARWATER
Ardenna bulleri
F: Procellariidae

This shearwater nests on islands off N. New Zealand but roams across the Pacific outside the breeding season.

sooty brown head

45–47 cm
18–18½ in

ANTARCTIC PRION
Pachyptila desolata
F: Procellariidae

Prions are small, grey Southern Ocean petrels that skim the sea to strain plankton with their flattened bill.




17–20 cm
6½–8 in



36–41 cm
14–16 in

SNOW PETREL
Pagodroma nivea
F: Procellariidae

One of the few birds that breed on the Antarctic, this petrel breeds further south than any other bird, even wandering to the South Pole.



41 cm
16 in

BLACK-CAPPED PETREL
Pterodroma hasitata
F: Procellariidae

Like many of the small, fast species known as gadfly petrels, this one has a tropical distribution. It breeds on islands of the West Indies.

39–40 cm
15½–16 in

CAPE PETREL
Daption capense
F: Procellariidae

One of several members of this family, mostly circumpolar in the southern hemisphere, the cape petrel breeds on islands around Antarctica and winters further north.



scattered dark feathers on white body

SOUTHERN GIANT PETREL
Macronectes giganteus
F: Procellariidae

This carrion-eating bird breeds in the South Atlantic. Unlike many other petrels, it has strong enough legs to walk well on land.

86–99 cm
34–39 in



massive yellowish bill

19–21 cm
7½–8½ in

BAND-RUMPED STORM PETREL
Oceanodroma castro
F: Hydrobatidae

Typical of northern hemisphere storm petrels, this tiny bird has a white-banded rump and forked tail. It is seen in both the Atlantic and Pacific.

