

**CATTLE EGRET***Bubulcus ibis*  
F: Ardeidae

Actually a small white heron, this globally widespread species forages on grassland and often follows livestock to catch disturbed prey.

48–53 cm  
19–21 in**INDIAN POND HERON***Ardeola grayii*  
F: Ardeidae

Common in S. Asia, this heron stalks aquatic prey, but it may also flush fish by flying low over water.

42–45 cm  
16½–18 in68–82 cm  
27–32 in

black-tipped bill

long neck plumes when breeding

grey body

**BOAT-BILLED HERON***Cochlearius cochlearius*  
F: Ardeidae

This heron has a broad bill, adapted for scooping animal prey. It inhabits C. and S. American mangrove swamps.

45–50 cm  
18–20 in**BLACK-CROWNED NIGHT HERON***Nycticorax nycticorax*  
F: Ardeidae

Night herons have good night vision. Found in most warm regions except Australia, this is the most widespread species.

58–65 cm  
23–26 in

bare head and neck

black wing plumes

65–75 cm  
26–30 in**STRAW-NECKED IBIS***Threskiornis spinicollis*  
F: Threskiornithidae

So called because of the straw-like feathers at the base of its neck, this nomadic ibis occurs in New Guinea and Australia.

59–76 cm  
23–30 in**SACRED IBIS***Threskiornis aethiopicus*  
F: Threskiornithidae

This is a common bird of the wetlands and grasslands of Africa and Madagascar. It has been introduced into America and Europe.

**SCARLET IBIS***Eudocimus ruber*  
F: Threskiornithidae


The national bird of Trinidad, the scarlet ibis occurs in tropical America and acquires its scarlet pigment from eating crustaceans.

56–61 cm  
22–24 in**REDDISH EGRET***Egretta rufescens*  
F: Ardeidae

This American species has white and reddish grey colour forms. When fishing, it sometimes holds out its wings to reduce sun glare.

**AUSTRALIAN IBIS***Threskiornis molucca*  
F: Threskiornithidae


This is a common species throughout Australia. It often invades urban areas, where it is sometimes considered a pest.

69–76 cm  
27–30 in76–89 cm  
30–35 in**HADADA IBIS***Bostrychia hagedash*  
F: Threskiornithidae

A common African species found in grasslands, forests, parks, and gardens, the hadada ibis is named after its distinctive flight call.

75–77 cm  
30 in**BLACK-FACED IBIS***Theristicus melanopsis*  
F: Threskiornithidae

Found in S. America, this species occurs in temperate grassland habitat from the Andes to Patagonia.

55–65 cm  
22–26 in**GLOSSY IBIS***Plegadis falcinellus*  
F: Threskiornithidae

Found throughout warm parts of the world, this is the most widespread ibis species. Its tree-nesting colonies may be accompanied by herons.