



BANDED STILT
Cladorhynchus leucocephalus
F: Recurvirostridae

Stilts hunt small swimming invertebrates. This Australian species gathers in large flocks on salt lakes to feed on brine shrimp.

36–45 cm
14–16 in



RED-NECKED AVOCET


Recurvirostra novaehollandiae
F: Recurvirostridae
A nomadic bird of Australian wetlands, this distinctively marked wader is a sociable species that feeds in large flocks.

40–46 cm
16–18 in

PIED AVOCET
Recurvirostra avosetta
F: Recurvirostridae
This Eurasian wader is typical of avocets: relatives of stilts that take small aquatic animals by sweeping their upturned bill through water.

42–45 cm
16½–18 in

black and white body



BLACK-WINGED STILT

Himantopus himantopus
F: Recurvirostridae
This highly variable stilt is nearly global in distribution. There are white-necked and black-necked forms that may be separate species.

33–36 cm
13–14 in



WRYBILL
Anarhynchus frontalis
F: Charadriidae

Related to lapwings, this wader from New Zealand is the only bird with a bill bent sideways, which it uses to reach invertebrates under stones.

20 cm
8 in



INLAND DOTTEREL
Peltohyas australis
F: Charadriidae

This sand-coloured relative of lapwings is found in arid parts of Australia, often far from water.

19–23 cm
7½–9 in



SPUR-WINGED PLOVER
Vanellus spinosus
F: Charadriidae

Occurring in wetlands across Africa and the Middle East, this bird is one of several lapwings, such as masked and southern, with a spur on each wing.

25–27 cm
10–10½ in



GREY PLOVER
Pluvialis squatarola
F: Charadriidae

The only member of the golden plover group with grey, not yellow, speckled upperparts, this species breeds around Arctic coastal tundra.

25–30 cm
10–12 in

black underparts when breeding




EURASIAN GOLDEN PLOVER
Pluvialis apricaria
F: Charadriidae

Golden plovers are more closely related to stilts and oystercatchers than they are to other plovers. This species develops a black belly when breeding.

26–29 cm
10–11½ in

blue-grey feet and legs



MASKED LAPWING
Vanellus miles
F: Charadriidae

Many lapwings have yellow, fleshy wattles. These facial adornments are especially prominent on this bird from Australasia.

35–38 cm
14–15 in



NORTHERN LAPWING
Vanellus vanellus
F: Charadriidae

Lapwings are named for the lapping sound of their flight. This Eurasian species has a distinctive, pointed crest. It is also called a peewit for its unique call.

28–31 cm
11–12 in



COMMON RINGED PLOVER
Charadrius hiaticula
F: Charadriidae

One of the most widespread ring-necked plovers, this species breeds around the Arctic and overwinters in Africa and S.W. Asia.



KILLDEER
Charadrius vociferus
F: Charadriidae

This long-tailed, grassland plover migrates between N. and S. America, but there are resident populations in Peru and Chile.

23–27 cm
9–10½ in



EURASIAN DOTTEREL
Charadrius morinellus
F: Charadriidae

Females of this tundra-breeding bird are more brightly coloured than males, but like many other plovers both become duller in winter.

20–22 cm
8–9 in