






LONG-BILLED CURLEW
Numenius americanus
F: Scolopacidae

A typical curlew, this American species is a large wader with a long, down-curved bill for probing deeply in mud for invertebrate prey.



WHIMBREL
Numenius phaeopus
F: Scolopacidae

A mid-sized curlew, the whimbrel has a distinctive trilling call and breeds in circumpolar regions. Some individuals winter as far away as Australia.



BUFF-BREASTED SANDPIPER
Tryngites subruficollis
F: Scolopacidae

This wader breeds in the tundra of N. America and extreme E. Siberia. It winters in the grasslands of S. America.



COMMON REDSHANK
Tringa totanus
F: Scolopacidae

Most waders breed near fresh water, but the Old World common redshank – named after its coloured legs – may sometimes breed on salt marshes.



LESSER YELLOWLEGS
Tringa flavipes
F: Scolopacidae

Yellowlegs belong to the shank group. This species breeds in forests in Alaska and Canada, and overwinters in the Caribbean.



WANDERING TATTLER
Tringa incana
F: Scolopacidae

This shank species breeds in Alaska, during which its breast is barred. It winters further south along the Pacific coastline of America.



AMERICAN WOODCOCK
Scolopax minor
F: Scolopacidae

Like other woodcocks and snipes, this American bird is superbly camouflaged and its eyes are placed high, enabling all-round vision for predators.



SPOTTED SANDPIPER
Actitis macularius
F: Scolopacidae

Like the related common sandpiper (*A. hypoleucos*) from Eurasia, this American sandpiper has a short bill for pecking food from dry ground.



RUDDY TURNSTONE
Arenaria interpres
F: Scolopacidae

This bird, which breeds across the northern hemisphere, is named for the way it flips objects to search for prey.



SPOON-BILLED SANDPIPER
Calidris pygmaea
F: Scolopacidae

Like spoonbills, this wader from E. Asia uses its distinctive spoon-like bill to sweep shallow water for invertebrate prey.



WHITE-BELLIED SEEDSNIPE
Attagis malouinus
F: Thinocoridae

Four species of short-billed, herbivorous seedsnipe live in the open habitats of S. America. This species is confined to the southern tip.



RUFF
Calidris pugnax
F: Scolopacidae

Males of this Old World marsh and meadow species transform during breeding: they lose their grey winter plumage and take on a rufous (reddish brown) and black coloration, with a showy neck ruff.

large neck ruff in breeding male

20–30 cm
8–12 in

18–20 cm
7–8 in

26–28 cm
10–11 in

26–30 cm
10–12 in

27–29 cm
10½–11½ in

23–25 cm
9–10 in

40–42 cm
16–16½ in

18–20 cm
7–8 in

45–66 cm
18–26 in

reddish legs