


» PIGEONS AND DOVES



33–40 cm
13–16 in

NICOBAR PIGEON
Caloenas nicobarica
F: Columbidae

Possibly related to the extinct Mauritian dodo, this pigeon inhabits coastal regions and island forests from Malaysia to New Guinea.



17–23 cm
6½–9 in

INCA DOVE
Columbina inca
F: Columbidae

From arid regions of the S. USA and C. America, this species belongs to a group of mostly drab-coloured tropical American ground-doves.




20 cm
8 in

DIAMOND DOVE
Geopelia cuneata
F: Columbidae

A small, nomadic bird from the dry interiors of Australia, this dove occurs in groups, flocking in large numbers at watering holes.

WOMPPOO FRUIT DOVE
Ptilinopus magnificus
F: Columbidae

Fruit doves are multicoloured relatives of imperial pigeons (*Ducula* sp.). This large species from New Guinea and Australia lives in the rainforest canopy.



29–55 cm
11½–22 in

yellow markings
on wings


dark green tail



25–31 cm
10–12 in

WHITE-TIPPED DOVE
Leptotila verreauxi
F: Columbidae

A tropical relative of the mourning dove, this species is widespread in C. and S. America, and reaches as far north as Texas.



30 cm
12 in

MINDANAO BLEEDING-HEART PIGEON
Gallicolumba criniger
F: Columbidae


The five species of Philippine bleeding-heart pigeons are named for their blood-red breast patch. This bird inhabits the southern islands of the archipelago.



35 cm
14 in

SULAWESI GROUND DOVE
Gallicolumba tristigmata
F: Columbidae

From the forests of Sulawesi, Indonesia, this terrestrial bird is related to Philippine bleeding-heart pigeons, and perhaps to various doves of arid Australia.



23–34 cm
9–13½ in

MOURNING DOVE
Zenaida macroura
F: Columbidae

So called because of its mournful call, this long-tailed dove occurs in open country throughout N. and C. America, and the Caribbean.



23–28 cm
9–11 in

EMERALD DOVE
Chalcophaps indica
F: Columbidae


An iridescent green bird, this ground-feeding species occurs in rainforests from India to islands of the southwest Pacific. It has a diet of fruits and seeds.



40–46 cm
16–18 in

TOPKNOT PIGEON
Lopholaimus antarcticus
F: Columbidae

A large hawk-like pigeon from E. Australia, this species is named for the double crest of feathers on its forecrown and crown.



45 cm
18 in

GREEN IMPERIAL PIGEON
Ducula aenea
F: Columbidae

Found in rainforest canopy, from India to S.E. Asia, this large bird has a deep, booming call and has a diet consisting mainly of fruits.



39–44 cm
15½–17½ in

PIED IMPERIAL PIGEON
Ducula bicolor
F: Columbidae

Imperial pigeons are large, fruit-eating rainforest birds. This species from S.E. Asia and Australasia has a white plumage that is often stained due to its diet.