

» CUCKOOS

26–32 cm
10–12½ in



YELLOW-BILLED CUCKOO

Coccyzus americanus
F: Cuculidae

One of several American tree cuckoos with brown plumage and white-spotted tails, this species migrates between N. and S. America.

38 cm
15 in



BLACK-BELLIED CUCKOO

Piaya melanogaster
F: Cuculidae

Like other American tree cuckoos, this species builds its own nest and cares for its young. It is found from Colombia to Bolivia.

GIANT COUA

Coua gigas
F: Cuculidae

Couas are Madagascan ground-cuckoos with blue facial skin and long eyelashes. This bird lives in dry coastal forests.

62 cm
24 in



34 cm
13½ in



GUIRA CUCKOO

Guira guira
F: Cuculidae

This shaggy-looking bird belongs to the ani group of S. American cuckoos. It occurs in noisy flocks and nests communally in trees.

48–52 cm
19–20½ in



GREATER COUCAL

Centropus sinensis
F: Cuculidae

Like other coucals, this S. Asian species is a strong-legged bird with spurred feet and long hind claws.

rufous
head crest

28 cm
11 in



PAVONINE CUCKOO

Dromococcyx pavoninus
F: Cuculidae

A smaller relative of the pheasant-cuckoo, this tropical S. American bird is a ground predator. It feeds on invertebrates.

long, graduated tail

36 cm
14 in



PHEASANT-CUCKOO

Dromococcyx phasianellus
F: Cuculidae

This American ground cuckoo inhabits the forest floor in tropical S. America, where it lays its eggs in the nests of smaller passerines.

streaked, reddish
brown wings



60–80 cm
23½–32 in

PHEASANT-COUCAL

Centropus phasianinus
F: Cuculidae

Parental care in coucals is provided mostly by the males. This Australasian species has a black body when breeding, and builds a cup-shaped nest in grass.

GROOVE-BILLED ANI

Crotophaga sulcirostris
F: Cuculidae

A heavy-billed American cuckoo occurring from California to Argentina, the communal-nesting ani is a clumsy flier but runs well.


white
cheek
stripe

grey-brown
upperparts with
white spots

33 cm
13 in



56 cm
22 in



GREATER ROADRUNNER

Geococcyx californianus
F: Cuculidae

This fast-running predator is an American ground cuckoo. The roadrunner inhabits deserts and nests in cacti, and occurs in a range from the USA to Mexico.

39–46 cm
15½–18 in

COMMON KOEL

Eudynamis scolopaceus
F: Cuculidae

Unusually for cuckoos, this tropical brood parasite, found from Asia to Australasia, is a fruit-eater. Males are black and females are grey-brown.

CORAL-BILLED GROUND CUCKOO

Carpococcyx renauldi
F: Cuculidae

One of three species of Asian ground cuckoos related to Madagascan couas, this bird inhabits rainforests of S.E. Asia.

65–68 cm
26–27 in

