

WHITE-EARED HUMMINGBIRD*Basilinna leucotis*

F: Trochilidae

This emerald hummingbird frequents pine and oak montane forests, often near streams, from S. Arizona to Nicaragua.

9–10 cm
3½–4 in

ANDEAN HILLSTAR*Oreotrochilus estella*

F: Trochilidae

An Andean coquette hummingbird, this species lives higher than other species of the group, and feeds in air so thin that it cannot hover.

straight, black bill

FORK-TAILED WOODNYMPH*Thalurania furcata*

F: Trochilidae

A relative of the C. American emerald hummingbirds, this S. American species occurs in lowland forests as far south as N. Argentina.

iridescent green throat

10 cm
4 in

13 cm
5 in

10 cm
4 in

9 cm
3½ in

LUCIFER HUMMINGBIRD*Calothorax lucifer*

F: Trochilidae

This bee hummingbird is found in a range from the southern USA to Mexico. It lives in semi-deserts, especially those with *Agave* plants.

purple throat patch

12 cm
4¾ in

BROAD-BILLED HUMMINGBIRD*Cynanthus latirostris*

F: Trochilidae

Males of this Mexican scrub hummingbird fly back and forth in a pendulum display to attract females. The species belongs to the “emerald” group.

23–26 cm
9–10 in

5–6 cm
2–2¼ in

SWORD-BILLED HUMMINGBIRD*Ensifera ensifera*

F: Trochilidae

The only bird with a bill longer than its body, this member of the Andean inca group favours the trumpet-shaped flowers of passion vines.

white spot near the eye

7–9 cm
2¾–3½ in

BEE HUMMINGBIRD*Mellisuga helenae*

F: Trochilidae

This tiny hummingbird is confined to Cuba, but is related to migratory N. American hummingbirds. Males of the species are the smallest of all birds.

RACKET-TAILED PUFFLEG*Ocreatus underwoodii*

F: Trochilidae

Belonging to the inca group, this bee-sized hummingbird of Andean forests is attracted to brush blossoms, such as those of legumes.

9 cm
3½ in

STRIPE-BREASTED STARthroat*Heliomaster squamosus*

F: Trochilidae

Male starthroats have a vividly coloured band around their throat. Although starthroats are related to C. American mountain gems, this species is from eastern Brazil.

12 cm
4¾ in

rufous (reddish brown) upperparts

9 cm
3½ in

RUFIOUS HUMMINGBIRD*Selasphorus rufus*

F: Trochilidae

Sustaining the longest migration of comparably small birds, this aggressively territorial bee hummingbird takes a route from Alaska to Mexico.

9 cm
3½ in

WHITE-NECKED JACOBIN*Florisuga mellivora*

F: Trochilidae

Genetic research suggests that this large, canopy-dwelling bird of tropical American lowlands belongs to a small lineage independent of other groups in its family.

GORGETED SUNANGEL*Heliangelus strophianus*

F: Trochilidae

Sunangels belong to the coquette group of Andean hummingbirds. This species is found from Colombia to Ecuador and lives in wet forest thickets.

CALLIOPE HUMMINGBIRD*Stellula calliope*

F: Trochilidae

One of the migratory bee hummingbirds, this bird breeds in open forests of western N. America and overwinters in the Mexican semi-desert.

PLOVERCREST*Stephanoxis lalandi*

F: Trochilidae

This distinctive bird may be related to larger sabrewings of C. America, but is found in montane forests of eastern S. America.