



beard-like bristles

22 cm  
9 in

**RED-FRONTED BARBET**  
*Tricholaema diademata*  
F: Ramphastidae

An E. African barbet related to the spot-flanked barbet, this species occurs in drier habitats. Like other barbets, it nests in tree-holes.



**TOUCAN-BARBET**  
*Semnornis ramphastinus*  
F: Ramphastidae

A bird of the rainforests of Colombia and Ecuador, this species is intermediate between barbets and toucans. It is exclusively a fruit-eater.

20 cm  
8 in22 cm  
9 in

**SPOT-FLANKED BARBET**  
*Tricholaema lacrymosa*  
F: Ramphastidae

A barbet of damp woodlands from C. to E. Africa, the spot-flanked barbet mainly feeds on figs and berries.

white-spotted  
upperparts15–16 cm  
6–6½ in

**D'ARNAUD'S BARBET**  
*Trachyphonus darnaudii*  
F: Ramphastidae

*Trachyphonus* barbets from Africa live in open country and spend much time on the ground. D'Arnaud's barbet occurs throughout E. Africa.

23 cm  
9 in

**RED-AND-YELLOW BARBET**  
*Trachyphonus erythrocephalus*  
F: Ramphastidae

A typical African ground barbet, this bird eats insects, fruits, seeds, and even small lizards. It often burrows into termite mounds to nest.

17 cm  
6½ in28 cm  
11 in


**FIRE-TUFTED BARBET**  
*Psilopogon pyrolophus*  
F: Ramphastidae

The only Asian barbet with a graduated tail, this S.E. Asian species also has tufts of facial bristles and a cicada-like call.

12–13 cm  
4¾–5 in

**GREEN-BACKED HONEYBIRD**  
*Prodotiscus zambesiae*  
F: Indicatoridae

Honeybirds eat insects, fruits, and even beeswax. This African species lays eggs in the nests of the white-eye, a small woodland bird, and its young kill host nestlings.




**NORTHERN WRYNECK**  
*Jynx torquilla*  
F: Picidae

This Eurasian woodland ant-eater has a weaker bill than true woodpeckers and is named for its ability to twist its neck.

**BAR-BREASTED PICULET**  
*Picumnus aurifrons*  
F: Picidae

Piculets are tiny nuthatch-like members of the woodpecker family. They use their short bill for taking insects from decayed wood. This species is from central S. America.

10 cm  
4 in10 cm  
4 in

**GOLDEN-SPANGLED PICULET**  
*Picumnus exilis*  
F: Picidae

This S. American piculet, like other piculets, lacks stiff, supporting tail feathers of bigger woodpeckers, so it spends less time on vertical trunks.

10 cm  
4 in

**OCHRACEOUS PICULET**  
*Picumnus limae*  
F: Picidae

Confined to E. Brazil, the ochraceous piculet – like other piculets – reuses woodpecker nesting holes, as its bill is too small to excavate its own nest.


10 cm  
4 in

**OCHRE-COLLARED PICULET**  
*Picumnus temminckii*  
F: Picidae

Confined to the forests of E. Paraguay, S.E. Brazil, and N.E. Argentina, this piculet is well camouflaged by its pattern and coloration.

**SPOTTED PICULET**  
*Picumnus pygmaeus*  
F: Picidae

The spotted piculet occurs only in the tropical forests of N.E. Brazil, where it is relatively common.

10 cm  
4 in