





NORTHERN FLICKER
Colaptes auratus
F: Picidae

American flickers are so called because of the way their coloured underwings flash in flight. This species has red- and yellow-shafted forms that hybridize.




HAIRY WOODPECKER
Picoides villosus
F: Picidae

A N. American relative of the three-toed woodpecker, this species becomes more abundant according to the availability of its prey – larvae of bark beetle.



GREAT SPOTTED WOODPECKER
Dendrocopos major
F: Picidae

A widespread member of a Eurasian group of pied woodpeckers, this bird is common in forests and gardens from Europe to S.E. Asia.



MIDDLE-SPOTTED WOODPECKER
Dendrocytes medius
F: Picidae

A pied woodpecker confined to Europe and S.W. Asia, this species drums less than the related great spotted woodpecker.




THREE-TOED JACAMAR
Jacamaralcyon tridactyla
F: Galbulidae

The most dull-coloured jacamar, this dry forest species nests in earth banks and has two toes in front and one behind.




GREAT JACAMAR
Jacamerops aureus
F: Galbulidae

This is the largest species of jacamar. It occurs from Costa Rica to Bolivia and feeds mainly on insects, supplemented by small lizards.



RUFIOUS-TAILED JACAMAR
Galbula ruficauda
F: Galbulidae

This C. and S. American species is typical of its genus, characterized by iridescent green coloration above and rufous below.



CHESTNUT JACAMAR
Galbalcyrhynchus purusianus
F: Galbulidae

Jacamars specialize in catching large insects, such as butterflies, from the air. This is one of two chestnut-coloured species from S. America.




BLACK-FRONTED NUNBIRD
Monasa nigrifrons
F: Bucconidae

Nunbirds are black-bodied relatives of puffbirds. This noisy S. American species often forages beneath monkey troops, feeding on disturbed small animal prey.



RUSTY-BREASTED NUNLET
Nonnula rubecula
F: Bucconidae

Nunlets are small, drab-coloured members of the puffbird family. This bird is a typical S. American species of vine-bordered forest.



WHITE-EARED PUFFBIRD
Nystalus chacuru
F: Bucconidae

Like other puffbirds, this species from central S. America is a big-headed, "puff-bodied" bird with a heavy bill for catching small animals.



SWALLOW-WINGED PUFFBIRD
Chelidoptera tenebrosa
F: Bucconidae

Described as martin-like when perched and bat-like in flight, this bird of northern S. America sallies for flying insects along rivers.

SERIEMAS

Seriemas, order Cariamiformes, are large, long-legged, noisy ground birds of dry savanna woodland in South America, with unclear relationships to other bird orders.

Once linked with cranes, seriemas have since been placed closer to falcons. The two species are among the largest ground-living birds of South America and rarely fly, except to reach the canopy of trees to nest and roost overnight. Their loud calls make them more often heard than seen.

PHYLUM	CHORDATA
CLASS	AVES
ORDER	CARIAMIFORMES
FAMILIES	1
SPECIES	2

loose plumage on neck and breast

barred wing feathers rarely revealed

RED-LEGGED SERIEMA
Cariama cristata
F: Cariamidae

Related to the extinct, predatory, giant terror birds, seriemas are natives of the grasslands of S. America. This species has a sickle-like claw to dismember small prey.

