

## MANAKINS

These birds, the Pipridae, are related to cotingas. The brightly coloured males gather in groups called leks to display to females. For many species, courtship involves elaborate dancing movements. Females build the nest and rear offspring alone.

15 cm  
6 in

### ARARIPE MANAKIN

*Antilophia bokermanni*

Described as recently as 1998, this critically endangered bird is found only in the Araripe highlands of N.E. Brazil.

11–13 cm  
4¼–5 in

### PIN-TAILED MANAKIN

*Ilicura militaris*

Males of this manakin from S.E. Brazil have elongated central tail feathers and display to females with raised rump feathers.

### BLUE MANAKIN

*Chiroxiphia caudata*

One of the most colourful birds of the rainforest of S. Brazil, the blue manakin has a whining, cat-like call.

14–15 cm  
5½–6 in

red cap

### STRIPED MANAKIN

*Machaeropterus regulus*

This is an inconspicuous species of northern S. America. Males of the species have been observed making insect-like buzzing sounds while displaying.

9–10 cm  
3½–4 in

### GOLDEN-HEADED MANAKIN

*Pipra erythrocephala*

The male of this S. American species displays by jumping and whirring its wings – and sliding to one side or backwards.

9 cm  
3½ in

## COTINGAS AND RELATIVES

A diverse group of tropical American passerines, the Cotingidae consists of fruit-eating or insect-eating forest birds.

Males are brightly coloured and are highly vocal when courting females; some species display in trees and others on the ground.

27–28 cm  
10½–11 in

### BARE-THROATED BELLBIRD

*Procnias nudicollis*

Bellbirds are named after their metallic calls. Like other bellbirds, males of this eastern S. American species have white plumage.

glossy black body

bright yellow throat

### FIERY-THROATED FRUIT-EATER

*Pipreola chlorolepidota*

Fruit-eaters are stocky green birds, males of which are usually black headed and red or yellow throated. This is a small Andean species.

13 cm  
5 in

bare red skin around eyes

22 cm  
9 in

### PURPLE-THROATED FRUITCROW

*Querula purpurata*

Males of this distinctive Amazonian cotinga have a prominent, iridescent purplish red throat patch. It can pluck fruits while hovering in mid-air.

28–30 cm  
11–12 in

purple-red patch

### BLACK-TAILED TITYRA

*Tityra cayana*

This canopy bird belongs to a group of big-headed fruit-eaters. It nests in tree-holes, often those abandoned by woodpeckers.

large crest hides the bill

brilliant red plumage

### ANDEAN COCK-OF-THE-ROCK

*Rupicola peruvianus*

The colourful crested males of this large Andean cotinga display in groups to females. Females nest among rocks and rear young alone.

28–32 cm  
11–12½ in