




GREAT KISKADEE
Pitangus sulphuratus
Widespread through tropical America and named after its call, this species is a typical sallying tyrant flycatcher, but also forages near the ground.

BOWERBIRDS AND CATBIRDS

The Australasian family Ptilonorhynchidae consists predominantly of fruit-eaters. Male bowerbirds – often brightly coloured – build structures called bowers to attract females. They mate with many females and play no part in raising the young.




GREEN CATBIRD
Ailuroedus crassirostris
Male catbirds – named for their cat-like call – attract mates by laying leaves on the ground. This species is found in New Guinea and E. Australia.



GOLDEN BOWERBIRD
Prionodura newtoniana
The male of this small species from N. Australia attracts a mate by building a tower of sticks up to 3 m (10ft) high.

AUSTRALASIAN TREECREEPERS

The birds of the family Climacteridae have evolved to resemble the northern treecreepers, but are unrelated and, unlike them, do not use their tail for support when climbing up trees.



BROWN TREECREEPER
Climacteris picumnus
This common eastern Australian species has a distinct dark-backed form in the north of the range and a brown-backed form in the south.

LYREBIRDS

The Menuridae are large Australian birds that eat ground insects. They have complex vocal organs and excel at mimicking forest sounds. Males court females on display mounds by fanning their long tails and plumes.



SUPERB LYREBIRD
Menura novaehollandiae
The commonest lyrebird, this species is found in forests of S.E. Australia and Tasmania. Its lyre-shaped outer tail feathers are adorned with notches.

OVENBIRDS AND RELATIVES

These American birds, the Furnariidae, are adept at hunting hidden invertebrates and are known for the immense diversity of their nest architecture. This includes stick nests, tunnel nests, and those that resemble clay ovens.



WHITE-EYED FOLIAGE-GLEANER
Automolus leucophthalmus
A S. American bird with distinctive white irises, like many insect-eaters, this species forages in mixed flocks to flush out prey.


RUFIOUS HORNERO
Furnarius rufus
Widespread in central and southern S. America, this species, typical of the hornero group, builds an oven-like mud nest.

HONEYEATERS

The birds of the family Meliphagidae are from Australia and the southwest Pacific islands. They feed on nectar with a long, brush-tipped tongue and are important pollinators of plants in that region. Other nectar-feeders, such as sunbirds, have evolved similar features.




SCARLET HONEYEATER
Myzomela sanguinolenta
This bird is an E. Australian member of a group of long-billed, flower-visiting honeyeaters that often carry daubs of pollen on their forehead.



BLUE-FACED HONEYEATER
Entomyzon cyanotis
A large, noisy honeyeater from Australia and New Guinea, this species is more insectivorous than many others, but it also feeds on fruits.



LEWIN'S HONEYEATER
Meliphaga lewinii
A member of a short-billed group of honeyeaters, this species from eastern Australia eats insects, fruits, and berries.



EASTERN SPINEBILL
Acanthorhynchus tenuirostris
Spinebills are an ancient group of heathland honeyeaters and the most specialized nectar-feeders of the family. This species is from E. Australia.




TUI
Prothemadera novaeseelandiae
Although confined to New Zealand, this species is related to short-billed honeyeaters of Australia. It has an extraordinary vocal range.



NEW HOLLAND HONEYEATER
Phylidonyris novaehollandiae
Like other honeyeaters, this white-whiskered species from southern Australia and Tasmania also takes honeydew – the waste sugar solution produced by certain sap-sucking insects.

WOODCREEPERS

The tropical American birds of the family Dendrocolaptidae are specialists at climbing up tree trunks. They have stiffened tails for support and strongly clawed front toes for gripping the bark.



SCALLOPED WOODCREEPER
Lepidocolaptes falcinellus
This typical buff and brown woodcreeper is found only in the forests of southeastern S. America.