





CUT-THROAT
Amadina fasciata

Named for the male's red neck patch, this is a common bird of dry African woodlands, and is often found near human habitations.



GOULDIAN FINCH
Erythrura gouldiae

A brilliantly coloured relative of parrotfinches, this endangered bird of northern Australia is nomadic. Males are red- or black-faced.

PIPITS AND WAGTAILS

Present on every continent, the birds of the family Motacillidae inhabit open country, where they feed on insects. Most wagtails have longer tails and are more brightly coloured than more sombre pipits, and some are associated with water.



GOLDEN PIPIT
Tmetothylacus tenellus
A bird of open bush and grassland, the golden pipit is confined to E. Africa, from Sudan to Tanzania.



RED-THROATED PIPIT
Anthus cervinus

This pipit breeds in Arctic tundra, during which it develops a coloured throat: reddish brown in males and pink in females.



BUFF-BELLIED PIPIT
Anthus rubescens

A typical ground-running pipit, this species breeds in Arctic tundra and winters in fields and on coasts further south.



YELLOW WAGTAIL
Motacilla flava

This widespread Eurasian species overwinters in Africa, India, and Australia. There are many races, including many with grey or black head patterns.



WHITE WAGTAIL
Motacilla alba

A typical wagtail, this species is widespread throughout Eurasia – often on farmland or in towns.


SUNBIRDS

The small, fast-moving, nectar-feeding Old World tropical birds of the family Nectarinidae are similar to American hummingbirds, with their long, decurved bill and long tongue. Males are usually brightly coloured with a metallic sheen. These birds are fiercely territorial.




STREAKY-BREASTED SPIDERHUNTER
Arachnothera affinis

Spiderhunters are drab, long-billed members of the sunbird family. Like other sunbirds, this S.E. Asian species eats invertebrates as well as nectar.



SCARLET-CHESTED SUNBIRD
Chalcomitra senegalensis

Common throughout much of sub-Saharan Africa, this large sunbird occupies a variety of wooded habitats.



PURPLE SUNBIRD
Cinnyris asiaticus

Like other sunbirds, this species from S. Asia feeds its young mostly on insects. The male loses its brilliant plumage after breeding.

WEAVERS

Gregarious seed-eaters, the Ploceidae build elaborate nests. Males usually take sole responsibility for this duty; females use the nest as the basis for choosing mates. Most species are African, with a few found in southern Asia.



CHESTNUT WEAVER
Ploceus rubiginosus

The genus *Ploceus* includes the largest number of species of weavers. This species occurs in E. Africa.



RED-BILLED QUELEA
Quelea quelea

Widely considered the world's most abundant bird, this African species gathers in giant flocks that can cause serious damage to crops.



RED-COLLARED WIDOWBIRD
Euplectes ardens

Breeding males of widowbirds are black and some fan enlarged tails during display flights. This species is widespread in sub-Saharan Africa.



YELLOW-CROWNED BISHOP
Euplectes afer

This African bird is related to widowbirds. Breeding males are vibrantly patterned; females and non-breeding males are red or black.

ACCENTORS

Mostly ground-dwelling passerines, the narrow-billed *Prunellidae* are found in Eurasia. Most species are adapted to high altitudes, but they move to lower altitudes in winter to supplement their insectivorous diet with seeds.

DUNNOCK
Prunella modularis

Unlike other accentors, this is a lowland bird, and it does not usually gather in flocks. It is widespread in temperate Eurasia.

