




BLUE-NAPED CHLOROPHONIA
Chlorophonia cyanea

The fruit-eating chlorophonias are mainly green birds. This species is widespread in forests across S. America.




PURPLE-THROATED EUPHONIA
Euphonia chlorotica

Euphonias are small birds that mainly eat fruits, especially mistletoe berries. This species is from northern S. America.



HOUSE FINCH
Carpodacus mexicanus

The N. American house finch is a type of rosefinch. Rosefinches are especially diverse in temperate Asia. The males are red or pale coloured.




PINE GROSBEEK
Pinicola enucleator

The pine grosbeak occurs in coniferous forests in the northern hemisphere. The bill of this grosbeak resembles that of the closely related bullfinches.




EURASIAN BULLFINCH
Pyrrhula pyrrhula

Bullfinches have a short, thick bill and a thick-set head. This is the most widespread of the bullfinches and is found in woodlands across temperate Eurasia.



IIWI
Drepanis coccinea

This long-billed nectar-eater is found only in Hawaii. It is a bird of montane forests, the largest populations occurring at higher elevations.



JAPANESE GROSBEEK
Eophona personata

A strikingly patterned bird, the Japanese grosbeak of cold northern forests breeds in Siberia and N. Japan and winters in S. China.

AMERICAN WARBLERS

Unrelated to warblers of the Old World, these finch-related insect-eaters of the family Parulidae are found throughout North and South America. The tropical species are sedentary, but temperate ones are migratory. The males moult their bright colours in winter.




BLACK-AND-WHITE WARBLER
Mniotilta varia

This N. American warbler moves up and down tree trunks like a nuthatch, and has long hind claws for clinging to bark.

YELLOW-BREASTED CHAT
Icteria virens

This N. American bird is relatively large for a warbler. It sings at night as well as during the day, and mimics the calls of other birds.



YELLOW WARBLER
Setophaga aestiva

Widespread from N. America to the Caribbean, there is a large number of local races of yellow warblers.



BAY-BREASTED WARBLER
Setophaga castanea

This bird breeds in spruce forests in eastern N. America. Its numbers fluctuate according to the abundance of its prey, the spruce budworm.



HOODED WARBLER
Setophaga citrina

Like the American redstart, this bird from broadleaf woodland of the E. USA feeds on insects by catching flies while in flight.



AMERICAN REDSTART
Setophaga ruticilla

An active forager, this N. American warbler habitually flashes its orange and black wings and tail to flush out insects. It also catches prey in the air.




NORTHERN WATERTHRUSH
Parkesia noveboracensis

A large, tail-bobbing ground warbler of N. America, this bird forages in leaf litter of wet woodlands and nests in thickets near water.




PROTHONOTARY WARBLER
Protonotaria citrea

Unusually for American warblers, this species of densely wooded swamps nests in tree cavities – sometimes using old woodpecker holes.



COMMON YELLOWTHROAT
Geothlypis trichas

A bird of wet habitats, this is a migratory species like other N. American warblers. It winters in California and Mexico.



GOLDEN-WINGED WARBLER
Vermivora chrysoptera

A warbler of eastern N. America, this bird breeds in scrubby, open habitats and farmland, and benefits from deforestation.