



35–50 cm
14–20 in

MARMOT

Marmota sp.


Several species of marmots inhabit mountain grasslands and rocky areas of N. America, while some occur in Eurasia. Marmots hibernate for up to eight months, depending on the species.

BLACK-TAILED PRAIRIE DOG

Cynomys ludovicianus

Active during daytime, this ground squirrel lives in large groups in “towns” consisting of extensive communal burrows.

21–27 cm
8½–10½ in



32–40 cm
12½–16 in


HARRIS'S ANTELOPE SQUIRREL

Ammospermophilus harrisi

An agile inhabitant of the Sonora Desert and northern Mexico, this rodent is active in the heat of the day, but hibernates during winter.

POCKET MICE AND KANGAROO RATS

Mostly common, with a few rare species, the Heteromyidae are found in varied habitats from Canada to Central America, with kangaroo rats living mainly in deserts. Pocket mice typically run on four feet, but kangaroo rats hop on two large hind feet.



7–9 cm
2¾–3½ in

DESERT POCKET MOUSE

Chaetodipus penicillatus

This species is one of many small nocturnal rodents that inhabit the open sandy deserts of the S.W. USA and N. Mexico.



10 cm
4 in


MERRIAM'S KANGAROO RAT

Dipodomys merriami

Bounding away with its tail outstretched, this nocturnal inhabitant of the N. American deserts looks like a miniature kangaroo.

JUMPING MICE

The families Dipodidae (jerboas), Zapodidae (jumping mice), and Sminthidae (birch mice) are made up of rodents that have a long tail and powerful hind feet that enable them to bound about like miniature kangaroos.



LESSER EGYPTIAN JERBOA

Jaculus jaculus

This desert species is found across N. Africa, from Senegal to Egypt, south to Somalia, and east to Iran.



MEADOW JUMPING MOUSE

Zapus hudsonius


Found in the cool grasslands of northern N. America, this species also has isolated populations in the mountains of Arizona and New Mexico.

7–11 cm
2¾–4¼ in

11–13 cm
4½–5 in

VOLES, LEMMINGS, AND MUSKRATS

About 765 chubby, short-tailed rodents, including hamsters, voles, lemmings, and muskrats, make up the family Cricetidae. They are found worldwide, from western Europe to Siberia and the Pacific coast.



EDIBLE DORMOUSE

Glis glis

Known in Germany as “the seven-sleeper”, this dormouse hibernates for over six months. It is eaten in Mediterranean countries, such as Slovenia.

13–19 cm
5–7½ in



COMMON DORMOUSE

Muscardinus avellanarius

An excellent climber and jumper, this dormouse feeds on flowers, fruits, and insects at night, in shrubby woodland across Europe.

6–9 cm
2¼–3½ in



9–12 cm
3½–4¾ in

COMMON VOLE

Microtus arvalis

A common burrowing inhabitant of grasslands across most of N. Europe, this vole also occurs further east and in Russia.



12–23 cm
4¾–9 in

EURASIAN WATER VOLE

Arvicola amphibius

In Britain and parts of Europe, this vole occurs near waterside habitats, but in Russia and Iran, it lives away from water, digging extensive burrow systems.



11–15 cm
4½–6 in

NORWAY LEMMING

Lemmus lemmus

The main small mammal of the European tundra, the lemming's fluctuating population affects the breeding success of many Arctic predators.



8–12 cm
3¼–4¾ in

STEPPE VOLE

Lagurus lagurus

The only species in its genus, this vole lives in the dry, grassy plains that stretch from Ukraine to W. Mongolia.



8–14 cm
3¼–5½ in

BANK VOLE

Myodes glareolus

Active mostly at dawn and night, this typical vole inhabits scrub, woodlands, and gardens across most of W. Europe and east into Russia.



25–30 cm
10–12 in

MUSKRAT

Ondatra zibethicus

Originally an inhabitant of rivers, ponds, and streams in N. America, the muskrat has been introduced to Europe and is now widespread.