


HOWLER, SPIDER, AND WOOLLY MONKEYS

The howler, spider, and woolly monkeys, and murequis of the Atelidae are the largest New World monkeys. All have a prehensile tail that is used as a fifth limb when moving through the trees. Spider monkeys have longer limbs than the other members of the family.

50–71 cm
20–28 in




GUATEMALAN BLACK HOWLER

Alouatta pigra

Found in the Yucatán Peninsula of Mexico, Belize, and Guatemala, this species forms groups of up to 11 individuals.

46–63 cm
18–25 in




MANTLED HOWLER

Alouatta palliata

Named after the “mantle” of long guard hair on its flanks, this howler monkey lives in C. America and northern S. America.

30–64 cm
12–25 in



COLOMBIAN BLACK SPIDER MONKEY

Ateles fusciceps rufiventris

Like other spider monkeys, this subspecies does not have thumbs on its hands. It is found in Colombia and Panama.

31–63 cm
12–25 in



GEOFFROY'S SPIDER MONKEY

Ateles geoffroyi

Active during the day, this fruit-eating species lives in relatively large groups of up to 35 individuals in forests throughout C. America.

46–78 cm
18–31 in



SOUTHERN MURIQUI

Brachyteles arachnoides

Also known as the woolly spider monkey, this species, found in Brazilian forests, is critically endangered due to the loss of its habitat.



hairless tail tip


45–65 cm
18–26 in

GREY WOOLLY MONKEY

Lagothrix cana

A powerfully built species, this monkey lives in large troops in the primary forests of Brazil, Bolivia, and Peru.

46–65 cm
18–26 in



HUMBOLDT'S WOOLLY MONKEY

Lagothrix lagotricha

One of the largest New World monkeys, this species lives in lowland primary forest in the upper Amazon basin.



48–63 cm
19–25 in

large larynx for howling

VENEZUELAN RED HOWLER

Alouatta seniculus

This howler monkey has an enlarged hyoid bone in the neck that allows it to make calls that can be heard several kilometres away.