



OLINGUITO
Bassaricyon neblina

Shy, nocturnal fruit-eaters, the olinguito is one of four species of olingo. It lives in the Andean cloud forests of Ecuador and Colombia.

30–40 cm
12–16 in



KINKAJOU
Potos flavus

A nocturnal tree-dweller from C. and S. America, the kinkajou uses its long tongue to pluck fruits and collect honey from bee nests.

41–76 cm
16–30 in

prehensile tail




NORTHERN RACCOON
Procyon lotor

This adaptable opportunist favours wooded and scrubby habitats across N. America, but often thrives in towns by scavenging human refuse.

long fur is grizzled

black “bandit mask” around eyes

44–62 cm
17½–24 in



RINGTAIL
Bassariscus astutus

An agile omnivore from C. America, this species forages by night for fruits and small animal prey, pausing often to scent-mark territory.

30–37 cm
12–14½ in



20–46 cm
8–18 in

EUROPEAN POLECAT
Mustela putorius

This lively nocturnal species, found in forests and meadows of C. and W. Europe, is the ancestor of the domestic ferret.



19–34 cm
7½–13½ in

STOAT
Mustela erminea

This lithe and ferocious little predator is found throughout much of the northern hemisphere. Its most northerly populations turn white in winter.



40–50 cm
16–20 in

BLACK-FOOTED FERRET
Mustela nigripes

Extinct in the wild in the late 20th century, this slender burrowing ferret has been reintroduced to reserves in the mid-western USA.




LEAST WEASEL
Mustela nivalis

The smallest carnivore, yet a fierce and highly successful predator, this weasel specializes in hunting mice.

11–26 cm
4¼–10 in

slender, elongated neck



20–26 cm
8–10 in

LONG-TAILED WEASEL
Mustela frenata

A widespread species of the Americas, this weasel hunts mice and voles. Northern individuals turn white in winter.