



AFRICAN ZORILLA
Ictonyx striatus

A striped polecat, this African species hunts a range of prey by night. It rests by day in hollow logs or burrows.

28–38 cm
11–15 in

long front
claws to dig up
buried insects

GIANT OTTER
Pteronura brasiliensis

This S. American predator needs 3 kg (2¼ lb) of fish a day to sustain itself. About 1,000–5,000 individuals of this endangered species remain.



1–1.3 m
3¼–4¼ ft



dorsal stripes meet
at the tail

24–33 cm
9½–13 in

**AFRICAN
STRIPED WEASEL**
Poecilogale albinucha

Found in C. and southern Africa, this species lives in self-dug burrows. It emerges at night to hunt for smaller animals, especially rodents, which it tracks by scent.

73–88 cm
29–35 in

AFRICAN CLAWLESS OTTER
Aonyx capensis

Found alongside water in forests and wetland throughout much of sub-Saharan Africa, this large otter eats mainly crabs, frogs, and fish.

36–47 cm
14–18½ in



greyish white
flecks on face
and throat

**ASIAN SMALL-CLAWED
OTTER**

Aonyx cinereus

The world's smallest otter, this species lives in wetlands in India and S.E. Asia. Its survival is threatened by habitat loss and pollution.


short,
blunt claws



50–82 cm
20–32 in

EURASIAN OTTER
Lutra lutra

Eurasian otters thrive in both river and coastal habitats, as long as they have access to fresh water for drinking and washing.



58–73 cm
23–29 in

NORTH AMERICAN RIVER OTTER
Lontra canadensis

Widespread in N. America, this species inhabits well-vegetated rivers and lake shores. It eats mainly fish and crayfish, but also hunts small land animals.



1–1.2 m
3¼–4 ft

SEA OTTER
Enhydra lutris

The sea otter hunts fish and shellfish in the cool North Pacific, relying on its incredibly dense fur to keep warm.