





PÈRE DAVID'S DEER
Elaphurus davidianus

Known only from captive populations, this deer was described in 1865 by a French missionary, Father Armand David, and has been reintroduced to China.

1.1–1.4 m
3½–4½ ft



AXIS DEER
Axis axis

Introduced in Australia and N. America, this common Indian forest deer has lyre-shaped antlers. It is a favourite prey of tigers.

70–95 cm
28–37 in




HOG DEER
Axis porcinus

Found in Asian forests, this deer is named after its hog-like habit of running head down, going under rather than leaping over obstacles.

55–75 cm
22–30 in

tail
white below



palmate
antlers

75–95 cm
30–37 in

long, pointed,
branched
antlers



BARASINGHA
Rucervus duvaucelii

This wetland deer from India was introduced in the USA for sport; the multi-pointed, branched antlers of males are highly prized.

1.2–1.4 m
4–4½ ft

50–70 cm
20–28 in




TUFTED DEER
Elaphodus cephalophus
This small deer lives in Asian montane forests. Males of this species have small antlers and short tusks, with a tuft of black hair on the forehead.

COMMON FALLOW DEER
Dama dama

Often domesticated as a source of venison, this species is distinguished by its spotted coat and flattened, palmate antlers.

0.95–1.3 m
3–4½ ft



WESTERN RED DEER
Cervus elaphus

From Europe, Turkey, and N. Africa, this deer varies across its range in bulk, antler size, and the prominence of its mane.

60–115 cm
23½–45 in



SIKA DEER
Cervus nippon

Distinguished by its stout, upright antlers, the sika breeds with red deer, especially when introduced outside its native E. Asia.

greyish brown
coat

WAPITI
Cervus canadensis

Although this N. American and Asian deer is similar in appearance to the red deer, genetic analysis has confirmed it is a species in its own right.

shaggy fur
on neck

75–105 cm
30–41¼ in



MULE DEER
Odocoileus hemionus

This species has a black-tipped tail and forked antlers, which distinguish it from the white-tailed deer that shares its range in western N. America.

55–105 cm
22–41¼ in



WHITE-TAILED DEER
Odocoileus virginianus

Found from Canada to Peru, and introduced in Europe and New Zealand, this deer flashes the white underside of its tail when alarmed.

reddish
brown coat

