

**TUFA**

This porous rock is formed by the precipitation of carbonate minerals from lower ambient temperature bodies of water than travertine, including those in cave systems.

layered structure

**GREY
ORTHOQUARTZITE****ORTHOQUARTZITE**

Rarely containing fossils, this rock is composed almost entirely of silica-cemented quartz grains. Orthoquartzite is sometimes known as quartz arenite.

**PINK
ORTHOQUARTZITE****MANGANESE NODULE**

Rich in valuable metals such as manganese, iron, nickel, cobalt, and copper, these nodules are rounded, black concretions that form on the deep ocean floor.

**LIGNITE****JET****LIGNITE**

Lignite is brown coal with a lower carbon content than bituminous coal. Jet is a black, lustrous form of lignite, and will take a high polish.

ANTHRACITE

The purest form of coal, which is derived from plant material, anthracite is black, with a vitreous lustre. It has curved edges when broken.

**BITUMINOUS COAL**

This sedimentary rock has a lower carbon content than anthracite. Brittle and dull, it is the most abundant type of coal.

*internal
radiating
structure*

*pale calcite
filling cracks
(septa)*

**SEPTARIAN
NODULE**

This type of concretion occurs in sedimentary rocks as individual, rounded masses. Internally, cracks known as septa are infilled by quartz or calcite.

**PYRITE NODULE**

These spherical nodules occur in shale and clay, and are composed entirely of radiating crystals of pyrite.

