



TABULATE CORAL
Catenipora sp.
A simple, tabulate coral with a chain-like structure, *Catenipora* lived in warm, shallow seas during Ordovician and Silurian times.

chain-like colonial structure



SCLERACTINIAN CORAL
Meandrina sp.
Shaped like the human brain, this colonial coral has ridges and valleys on its surface. First found in Eocene rocks, *Meandrina* still survives today.



RUGOSE CORAL
Goniophyllum pyramidale
A solitary coral occurring in Silurian rocks, *Goniophyllum* has a cone-shaped structure in which the polyp lived.

thick corallite wall

CAMBRIAN TRILOBITE
Paradoxides bohemicus
Some *Paradoxides* trilobites grew to nearly 1 m (3¼ ft) in length. This species had long spines on its body and is from Cambrian strata.



ORDOVICIAN TRILOBITE
Eodalmantina macrophtalma
An Ordovician trilobite, *Eodalmantina* had large, crescent-shaped eyes. Its thorax was composed of 11 segments.


SILURIAN TRILOBITE
Dalmanites caudatus
Common in the Silurian Period, *Dalmanites* had a segmented thorax and a pointed tail spine.



DEVONIAN ENROLLED TRILOBITE
Phacops sp.
Characterized by compound eyes, *Phacops* is found in Devonian strata. Trilobites could roll up, like many modern arthropods.



HORSESHOE CRAB RELATIVE
Euproops rotundatus
A Carboniferous relative of horseshoe crabs, *Euproops* had a crescent-shaped headshield and long tail spine.



LOBSTER
Eryma leptodactylina
A fossil lobster from rocks of Jurassic and Cretaceous times, *Eryma* was 6 cm (2¼ in) long, and similar to modern species.

pincers



CRAB
Avitelmessus grapsoides
Covered with many spines, this crab is from the Cretaceous Period. It grew to 25 cm (10 in) in width.



COCKROACH RELATIVE
Archimylacris eggintoni
A relative of cockroaches from the Carboniferous Period, *Archimylacris* had hind wings with a distinctive vein pattern.